

## **RESEARCH ARTICLE**

### **SCREENING OF ETHIOPIAN FABA BEAN (*VICIA FABA* L.) ACCESSIONS FOR RESISTANCE AND YIELD PARAMETERS AGAINST FABA BEAN GALL DISEASE (*OLPIDIUM VICIAE*) UNDER FIELD CONDITIONS**

Alemayehu Dugassa<sup>1</sup>, Tesfaye Alemu<sup>1,\*</sup>, Yitbarek Woldehawariat<sup>2</sup>, Serawit Handiso<sup>3</sup>, and Gezahegne Getaneh<sup>4</sup>

**ABSTRACT:** Faba bean (*Vicia faba* L.) has been produced for centuries in Ethiopia and provides protein supplements to the diet of rural households. It is recently threatened by a new emerging gall disease caused by the fungal pathogen *Olpidium viciae* Kusano. The purpose of this study was to screen faba bean accessions with high yield potential for resistance against gall disease under rain-fed field conditions. A local check and 180 accessions were screened following standardized procedures. The seeds were sown in a randomized complete block design with three replications during the 2017–2019 cropping seasons. SPSS statistical software version 20 was used to calculate the disease severity index (DSI), an area under disease progress curve. Significant ( $p \leq 0.05$ ) positive correlation was recorded between the disease severity and altitude ( $r = 0.27$ ), precipitation ( $r = 0.37$ ). Though no accessions were found resistant, 87.18% were identified as moderately resistant (DSI: 16.50–39.45%) to the test pathogen. In a combined mean analysis over districts, the lowest DSI was recorded for FB-Obse (DSI=16.50%), FB-Hachalu (DSI = 17.83%), FB-26872 (DSI = 18.84%), FB-26873 (DSI = 9.61%), and FB-28107 (DSI = 19.85%) accessions. Similarly, they exhibited various levels of yield potentials. The highest DSI rate (74.88–100%) was recorded between 38–77 days after crop emergence. The average minimum infection rate was recorded ( $r = 0.003$ ) for FB-26873 variety at Sululta with a coefficient of determination of 79.00%. Crops with the lowest DSI exhibited the uppermost yield parameters. Hence, faba bean accessions with the maximum yield and moderate resistance to gall disease were recommended to be used for production.

**Key words/phrases:** Disease severity index, Faba bean, Gall disease, Resistant, Yield parameter.

---

<sup>1</sup> Department of Microbial, Cellular and Molecular Biology, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia. E-mail: tesfaye.alemu@aau.edu.et

<sup>2</sup> Department of Zoological Sciences, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia.

<sup>3</sup> Department of Plant Sciences, College of Agriculture, Wolaita Sodo University

<sup>4</sup> Ambo Plant Protection Research Centre, Ethiopian Institute of Agricultural Research, Ethiopia.

\*Author to whom all correspondence should be addressed

## INTRODUCTION

Faba bean (*Vicia faba* L.) is an important pulse crop cultivated in the world for food and fodder (Singh *et al.*, 2013). Ethiopia is the main faba bean diversity centre next to China (Behailu Mulugeta *et al.*, 2018). Faba bean is mostly grown by farmers in the highlands of Ethiopia (1,800–3,000 m.a.s.l) receiving annual rainfall of 700–1,000 mm (Teklay Abebe *et al.*, 2018). It has an important place in the Ethiopian national diet and is used to make various traditional dishes for its high protein content (Endalkachew Fekadu *et al.*, 2018; Etemadi *et al.*, 2019).

Despite its wide ecological and economic benefits, the average actual yield (2.1 tons per hectare) of faba bean is far below its potential 4.4 tons per hectare (Wondafrash Mulugeta *et al.*, 2019). The low crop yield is attributed to different biotic and abiotic factors (Alireza *et al.*, 2021). Among biotic factors, fungal disease such as chocolate spot (caused by *Botrytis fabae* Sardiña), rust (caused by *Uromyces viciae-fabae* (Pers.) J. Schröt.), downy mildew (caused by *Peronospora viciae* (Berk.) Casp.), and black root rot (caused by *Fusarium solani* (Mart.) Sacc.) are the most common widespread and destructive among faba bean yield-limiting fungal diseases (Teklay Abebe *et al.*, 2014). Similarly, faba bean productivity is significantly influenced by its genotype (Sheikh *et al.*, 2015).

In recent years, in addition to the previous common diseases, the crop is threatened by the new emerging faba bean gall disease caused by *Olpidium viciae* Kusano. The gall disease was widely disseminated in the high altitudes (above 2,400 m a.s.l.) of faba bean growing areas within a few years since 2010 (Teklay Abebe *et al.*, 2018). The disease becomes the most devastating and causes up to 100% crop loss in many highland areas of the country (Bogale Nigir *et al.*, 2017).

Although farmers use different management options such as good soil drainage, use of disease-free seeds, fungicide spray, and crop rotation, none of these were able to control gall disease adequately under field conditions (Teklay Abebe *et al.*, 2018). Fungicides have negative effects on public health and environmental balance (Singh *et al.*, 2013). Hence, it needs more attention to find non-hazardous solutions to manage the disease and sustain the productivity of the crop.

In Ethiopia, it is clearly observed that there is no sound gall disease control measure identified yet. Screening of faba bean accessions being resistant to gall disease and having higher productivity potential could be the most sustainable management options. The cultivation of resistant crop varieties

eliminates the crop losses caused by disease (Soler *et al.*, 2021; Su *et al.*, 2021) and reducing costs related to fungicide applications, and increases crop productivity (Pornthip and Prakrit, 2021). The hazard of pollution of the environment with toxic chemical compounds can also be avoided (Teklay Abebe *et al.*, 2014).

As a result of efforts made through national and regional faba bean improvement programs of Ethiopia, many faba bean varieties were released at different times. There are only a few studies about the evaluation of resistance of faba bean varieties against gall diseases in Ethiopia. Therefore, the objective of this study was to evaluate and identify nationally available faba bean accessions for high yield and better resistance to faba bean gall diseases after natural infection at the most faba bean gall disease hot spot areas under rain-fed field conditions.

## MATERIALS AND METHODS

### Description of the study sites

The study was conducted at Angolelana Tera, Sululta, and Midakegn districts on farmers' fields during the main cropping seasons (June–August) of the year 2017–2019. These study sites are considered the hotspot areas for faba bean gall disease since its occurrence in Ethiopia (Bitew Beyene and Tigabie Abiro, 2016). Angolelana Tera is located in North Shoa zone of the Amhara National Regional State, and 110 km North East of the capital city, Addis Ababa. It is found at the longitude of 39° 27' 0" E and latitude of 9° 33' 0" N. The altitude of the specific study site is 2,774 m.a.s.l. with brown soil colour and plane topography.

Both Sululta and Midakegn districts are located in the Oromia National Regional State. Sululta district is located in North Shoa zone at about 23 km to the North of Addis Ababa. It is found at longitude 38° 43' 59.99" E and latitude 9° 10' 59.99" N, and an altitude of the specific study site is 2,610 m.a.s.l., whereas Midakegn district is located in West Shoa zone at about 170 km to the West of Addis Ababa, Ethiopia. It is found at longitude 37° 28' 94" E, latitude 9° 08' 61" N and altitude 2520 m.a.s.l. Soil types of all the research sites are vertisols.

In the 2017 cropping season, the monthly mean precipitation for June–August was 284.40 mm, 273.90 mm, and 231 mm at Angolelana Tera, Sululta, and Midakegn districts, respectively. Whereas the average daily minimum mean temperature (14.60°C) was recorded at Midakegn district. Similarly, in the 2018 cropping season, the maximum monthly mean

precipitation (358.20 mm) was recorded at Angolelana Tera district. During this period the average temperature at all sites was between 15.40–15.50°C. Likewise, in the 2019 cropping season, the maximum precipitation was recorded at Angolelana Tera (274.50 mm) followed by Midakegn district (272.7 mm), whereas the minimum average daily temperature was recorded at Angolelana Tera district (14.90°C) (Table 1).

Table 1. Climate data of the month June to August during 2017–2019 cropping seasons.

| Districts and cropping seasons  | Precipitation (mm)<br>Mean ± SD | Daily temperature (°C) |              |              |
|---------------------------------|---------------------------------|------------------------|--------------|--------------|
|                                 |                                 | Minimum                | Maximum      | Average      |
|                                 |                                 | Mean ± SD              | Mean ± SD    | Mean ± SD    |
| <b>Angolelana Tera district</b> |                                 |                        |              |              |
| 2017                            | 284.40 ± 90.40                  | 8.60 ± 1.10            | 21.20 ± 0.90 | 14.90 ± 0.20 |
| 2018                            | 358.20 ± 50.10                  | 9.30 ± 0.90            | 21.50 ± 0.50 | 15.40 ± 0.70 |
| 2019                            | 274.50 ± 64.90                  | 8.60 ± 0.60            | 21.30 ± 0.90 | 14.90 ± 0.50 |
| Average                         | 295.70 ± 78.90                  | 8.80 ± 0.80            | 21.30 ± 0.70 | 15.10 ± 0.50 |
| <b>Sululta district</b>         |                                 |                        |              |              |
| 2017                            | 273.90 ± 91.30                  | 9.60 ± 0.90            | 22.50 ± 0.90 | 16.10 ± 0.10 |
| 2018                            | 349.10 ± 53.20                  | 9.20 ± 0.60            | 21.90 ± 0.10 | 15.50 ± 0.30 |
| 2019                            | 206.90 ± 86.40                  | 9.00 ± 0.10            | 22.10 ± 0.60 | 15.60 ± 0.30 |
| Average                         | 276.70 ± 91.90                  | 9.30 ± 0.70            | 22.20 ± 0.60 | 15.70 ± 0.30 |
| <b>Midakegn district</b>        |                                 |                        |              |              |
| 2017                            | 231.00 ± 75.50                  | 10.00 ± 1.20           | 19.30 ± 0.60 | 14.60 ± 1.00 |
| 2018                            | 277.70 ± 31.50                  | 9.70 ± 0.20            | 21.10 ± 0.30 | 15.40 ± 0.20 |
| 2019                            | 272.70 ± 58.50                  | 9.90 ± 0.70            | 21.20 ± 0.70 | 15.50 ± 0.20 |
| Average                         | 240.90 ± 55.70                  | 9.90 ± 0.20            | 19.80 ± 1.60 | 15.90 ± 1.20 |

Source: EMA, 2019

## Collection of faba bean accessions and screening procedures

A total of 180 faba bean accessions were collected from the Ethiopian Biodiversity Institute (EBI), and Holota Agricultural Research Centre (HARC). The screening processes of faba bean accessions against gall diseases were conducted for three consecutive cropping seasons from 2017 to 2019 under natural infection in rain-fed field conditions. The faba bean germplasm were sown in June (main rainy season) for all years. In the 2017 cropping season, all the collected accessions were sown in the farmers' fields at each experimental site and those best adapted to the study locations (30 from EBI and eight from HARC collection) were selected for additional screening in the 2018 cropping season. Based on the results of the 2017 and 2018 cropping season, a local check and 14 faba bean accessions with moderate resistance against gall diseases and potential yield parameters were selected and planted for further evaluation in the 2019 cropping season.

## Experimental design and field management

Farmland previously sown with faba bean was prepared with assistance from the farm field owners (farmers) for the three cropping seasons (2017–2019) at each experimental site. In all cropping seasons, the faba bean accessions were arranged in Randomized Complete Block Design (RCBD) with three replications. In 2017, a total of 180 faba bean accessions were sown on a 500 m<sup>2</sup> (25 m length x 20 m width) experimental field area which was divided into ten blocks. Each block consisted of 24 plots. The faba bean accessions were evaluated for gall disease resistance and adaptation to the ecological zone of the study areas. Then, 38 faba bean accessions that showed better adaptation to the study area and tolerant to gall disease were screened for the 2018 cropping season. In 2018, a total of 200 m<sup>2</sup> (20 m length x 10 m width) experimental farm fields were prepared and divided into six blocks. Each block consisted of 16 plots. At the final experimental period (2019 cropping season) a total of 100 m<sup>2</sup> (10 m length x 10 m width) experimental field was divided into four blocks. Each block consisted of ten (10) plots. Each block was 40 cm far apart in all experimental periods. Similarly, each plot was arranged in 1 m<sup>2</sup> in area and divided into three rows. The spaces between rows were 40 cm. A total of 12 seeds were sown for each accession (4 seeds/row) on each plot. The distance between seeds was 10 cm. The local check was sown in every 10 test plots. The experimental field was plowed following the standard practices for faba bean production before sowing. Weeding and other agronomic practices were carried out at an appropriate time.

## Data collection

Disease and crop data were recorded on plant basis to evaluate the performance of the experimental accessions in different time intervals. The disease data were recorded on the 60th day after the date of crop emergence. Percent disease incidence of faba bean gall diseases was assessed by counting the number of diseased plants per total number of plants inspected and expressed as a percentage of total plants. The gall disease severity index was recorded using a 0–9 scale following the methods described by Aghajani (2009). Six resistance levels were used in these experiments viz., highly resistant (healthy plant, grade = 0, DSI = 0–2%), resistant (1–15 % of the plant surface infected, grade = 1, DSI = 2.1–15%), moderately resistant (15.1–40% of the plant surface infected, grade = 3, DSI = 15.1–40 %), moderately susceptible (40.1–60 of the plant surface infected, grade = 5, DSI = 40.1–60%), susceptible (60.1–80% of the plant surface infected,

grade = 7, DSI = 60.1–80%), highly susceptible (>80% of plant surface infected, grade = 9, DSI = 80.1–100%). The DSI of each faba bean accession was calculated by the following formula (Aghajani, 2009).

$$DSI = \left( \frac{a + 3b + 5c + 7d + 9e}{9(a + b + c + d + e)} \right) \times 100$$

where, 9 = highest rating value, a = number of plants in class 1, b = number of plants in class 3, c = number of plants in class 5, d = number of plants in class 7, e = number of plants in class 9.

The Area Under the Disease Progress Curve (AUDPC) was calculated from DSI (Villegas-Fernandez *et al.*, 2011) as follows;

$$AUDPC = \sum_{i=0}^n [0.5 (y_i + y_{i+1}) (t_{i+1} - t_i)]$$

Where  $y_i$  is disease severity at  $i^{th}$  assessment,  $t_i$  is a time of assessment and  $n$  is total assessments.

The crop was harvested 115 days after the date of crop emergence. The yield parameters (viz., pod/plant, seed/pod, and 100 seeds weight) of each faba bean variety were evaluated for each cropping season and experimental sites. Hundred seed weight was determined using a digital balance (Sartorius, Germany).

### Data analysis

The data were analysed by one-way ANOVA implemented in SPSS statistical software version 20 to evaluate the presence of significant difference among faba bean accessions in response to gall disease. The AUDPC of the disease was analyzed from DSI. Means were compared using the least significant difference (LSD) at a 5% probability level ( $p \leq 0.05$ ). The percentage of seed germination in the field was calculated by counting the number of seeds germinated per total number of seed sown and expressed as a percentage of total seeds (Bhagat *et al.*, 2013).

## RESULTS

### Gall disease parameters

#### Gall disease intensity in 2017 and 2018 cropping seasons

The experiment results revealed that there was significant variation ( $p < 0.05$ ) among the faba bean accessions against gall disease incidence and the disease severity index (DSI) in combined mean analysis over the experimental sites, and cropping seasons. FB-28104 accession was found to have the highest percent disease incidence with 94.22%. This was followed by a statistically non-significant ( $p > 0.05$ ) result of FB-212566 (93.39%) (Table 2).

The combined mean DSI of faba bean gall-forming diseases in 2017 and 2018 cropping seasons were between 15.60% and 61.50%. The highest and the lowest disease DSI were recorded on FB-26869 (61.50%) and FB-26872 (15.60%) accessions, respectively. There was significant variability ( $p \leq 0.05$ ) in the resistance of faba bean accession against faba bean gall disease (Table 2). In this study, the average of two years' experiments indicated that among 38 accessions, and a local check, 34 (87.18%) were classified as moderately resistant (DSI: 15.60–39.45%), four (10.26%) were classified as moderately susceptible (DSI: 41.35–41.47%), and one (2.56%) was susceptible (FB-26869: DSI = 61.50%). However, no resistant accession was found against faba bean gall disease. Hence, fourteen faba bean accessions (viz., FB-26872, FB-Tumsa, FB-26873, FB-28107, FB-Obse, FB-Hachalu, FB-26883, FB-Numan, FB-28770, FB-26885, FB-COLL-0038, FB-26875, FB-26867, and FB-25280) those had shown better reactions among moderately resistant accessions were selected for further evaluation (Table 2).

### **Gall disease intensity during 2019 cropping season**

The result presented in Table 3 indicated that the reaction of 14 faba bean accessions and a local check against faba bean gall disease was significantly varied ( $p \leq 0.05$ ) across districts. The lowest DSI was recorded on FB-Hachalu (6.67%) followed by FB-Obse (8.33%) at Sululta district than the remaining accessions. Similarly, the lowest mean DSI of gall disease was recorded on FB-Obse accession with 11.67%, and 16.33% at Midakegn, and Angolelana Tera districts, respectively. The highest DSI was recorded on local check at Angolelana Tera (DSI = 63.00%) and Sululta (DSI = 22.38%) districts, whereas the highest DSI recorded on FB-26885 (DSI = 46.67%) accession at Midakegn district (Table 3).

In the analysis of a combined mean average faba bean accessions, the lowest average DSI (13.11%) was recorded at Sululta, whereas the higher average DSI (38.76%) was recorded at Angolelana Tera than Midakegn district (DSI = 31.89%) (Table 3).

Table 2. Average faba bean gall diseases intensity of the three experimental sites during 2017 and 2018 cropping seasons of some faba bean accessions.

| Faba bean accessions | 2017 cropping season       |                            | 2018 cropping season       |                            | Combined mean              |                            | Reaction |
|----------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------|
|                      | Incidence (%)<br>± SD      | Severity (%)<br>± SD       | Incidence (%) ±<br>SD      | Severity (%)<br>± SD       | Incidence (%)<br>± SD      | Severity (%)<br>± SD       |          |
| FB-26872             | 88.83 <sup>b</sup> ± 3.00  | 16.83 <sup>c</sup> ± 0.35  | 50.79 <sup>a</sup> ± 42.07 | 14.36 <sup>a</sup> ± 4.90  | 69.08 <sup>a</sup> ± 26.90 | 15.60 <sup>a</sup> ± 1.74  | MR       |
| FB-Tumsa             | 80.00 <sup>b</sup> ± 11.27 | 20.15 <sup>a</sup> ± 10.56 | 52.47 <sup>a</sup> ± 40.73 | 13.17 <sup>a</sup> ± 1.52  | 66.24 <sup>a</sup> ± 19.46 | 16.66 <sup>a</sup> ± 9.94  | MR       |
| FB-26873             | 91.20 <sup>b</sup> ± 8.32  | 17.04 <sup>c</sup> ± 5.13  | 58.32 <sup>c</sup> ± 36.72 | 16.61 <sup>a</sup> ± 4.81  | 74.76 <sup>b</sup> ± 32.40 | 16.83 <sup>a</sup> ± 0.30  | MR       |
| FB-28107             | 97.17 <sup>a</sup> ± 12.10 | 21.08 <sup>a</sup> ± 1.25  | 60.31 <sup>c</sup> ± 34.11 | 15.94 <sup>a</sup> ± 3.83  | 78.74 <sup>b</sup> ± 26.06 | 18.51 <sup>a</sup> ± 3.63  | MR       |
| FB-Obse              | 98.60 <sup>a</sup> ± 11.23 | 21.05 <sup>a</sup> ± 1.23  | 53.32 <sup>a</sup> ± 39.23 | 16.34 <sup>a</sup> ± 3.67  | 75.96 <sup>b</sup> ± 32.02 | 18.70 <sup>a</sup> ± 3.33  | MR       |
| FB-Hachalu           | 97.67 <sup>a</sup> ± 10.45 | 21.23 <sup>a</sup> ± 0.45  | 51.85 <sup>a</sup> ± 44.93 | 16.15 <sup>a</sup> ± 3.63  | 74.76 <sup>b</sup> ± 32.40 | 18.79 <sup>a</sup> ± 3.59  | MR       |
| FB-26883             | 91.14 <sup>b</sup> ± 21.25 | 18.21 <sup>c</sup> ± 3.56  | 60.35 <sup>c</sup> ± 32.80 | 20.87 <sup>b</sup> ± 11.77 | 75.75 <sup>b</sup> ± 21.77 | 19.54 <sup>a</sup> ± 1.88  | MR       |
| FB-Numan             | 90.17 <sup>a</sup> ± 10.25 | 21.54 <sup>a</sup> ± 2.53  | 59.45 <sup>c</sup> ± 33.94 | 16.46 <sup>a</sup> ± 3.96  | 74.81 <sup>b</sup> ± 21.72 | 19.00 <sup>a</sup> ± 3.59  | MR       |
| FB-28770             | 85.00 <sup>b</sup> ± 11.25 | 22.44 <sup>a</sup> ± 2.56  | 56.32 <sup>c</sup> ± 37.41 | 18.44 <sup>a</sup> ± 6.01  | 70.66 <sup>a</sup> ± 20.28 | 20.44 <sup>a</sup> ± 2.83  | MR       |
| FB-26885             | 98.83 <sup>a</sup> ± 12.45 | 21.75 <sup>a</sup> ± 4.05  | 57.42 <sup>c</sup> ± 36.58 | 20.92 <sup>b</sup> ± 6.67  | 78.13 <sup>b</sup> ± 29.28 | 21.33 <sup>a</sup> ± 0.59  | MR       |
| FB-COLL-0038         | 96.80 <sup>a</sup> ± 18.45 | 21.05 <sup>a</sup> ± 0.45  | 71.57 <sup>b</sup> ± 27.10 | 21.01 <sup>b</sup> ± 9.00  | 84.19 <sup>c</sup> ± 17.84 | 21.03 <sup>a</sup> ± 0.03  | MR       |
| FB-26875             | 95.20 <sup>a</sup> ± 9.20  | 21.60 <sup>a</sup> ± 0.76  | 64.66 <sup>c</sup> ± 31.15 | 21.09 <sup>b</sup> ± 9.09  | 79.93 <sup>b</sup> ± 21.60 | 21.34 <sup>a</sup> ± 0.36  | MR       |
| FB-26867             | 91.00 <sup>b</sup> ± 6.02  | 23.67 <sup>b</sup> ± 1.45  | 46.03 <sup>d</sup> ± 45.78 | 21.13 <sup>b</sup> ± 11.58 | 68.52 <sup>a</sup> ± 31.80 | 22.40 <sup>a</sup> ± 1.79  | MR       |
| FB-25280             | 95.00 <sup>a</sup> ± 0.98  | 17.83 <sup>c</sup> ± 2.47  | 53.73 <sup>a</sup> ± 43.53 | 28.24 <sup>c</sup> ± 27.72 | 74.36 <sup>b</sup> ± 29.18 | 23.04 <sup>a</sup> ± 7.36  | MR       |
| FB-25018             | 94.17 <sup>a</sup> ± 1.23  | 21.25 <sup>a</sup> ± 3.12  | 69.95 <sup>b</sup> ± 25.26 | 41.77 <sup>d</sup> ± 20.96 | 82.06 <sup>c</sup> ± 17.13 | 31.51 <sup>b</sup> ± 14.51 | MR       |
| FB-26871             | 98.60 <sup>a</sup> ± 8.69  | 21.60 <sup>a</sup> ± 5.02  | 73.42 <sup>b</sup> ± 23.92 | 42.22 <sup>d</sup> ± 11.59 | 86.01 <sup>c</sup> ± 17.80 | 31.91 <sup>b</sup> ± 14.58 | MR       |
| FB-28112             | 86.33 <sup>b</sup> ± 12.24 | 22.67 <sup>a</sup> ± 0.25  | 64.73 <sup>c</sup> ± 31.49 | 42.92 <sup>d</sup> ± 12.75 | 75.53 <sup>b</sup> ± 15.28 | 32.80 <sup>b</sup> ± 14.32 | MR       |
| FB-212565            | 98.17 <sup>a</sup> ± 23.58 | 24.75 <sup>b</sup> ± 0.14  | 74.45 <sup>b</sup> ± 18.71 | 43.49 <sup>d</sup> ± 14.15 | 86.31 <sup>c</sup> ± 16.77 | 34.12 <sup>b</sup> ± 13.25 | MR       |
| FB-Dosha             | 99.00 <sup>a</sup> ± 20.12 | 24.10 <sup>b</sup> ± 3.54  | 64.06 <sup>c</sup> ± 29.64 | 44.59 <sup>d</sup> ± 13.43 | 81.53 <sup>c</sup> ± 24.71 | 34.35 <sup>b</sup> ± 14.49 | MR       |
| FB-26870             | 98.33 <sup>a</sup> ± 10.25 | 22.67 <sup>a</sup> ± 0.25  | 76.82 <sup>f</sup> ± 29.80 | 47.46 <sup>e</sup> ± 14.78 | 87.58 <sup>c</sup> ± 15.21 | 35.07 <sup>b</sup> ± 17.53 | MR       |
| FB-26400             | 85.67 <sup>b</sup> ± 11.25 | 23.25 <sup>a</sup> ± 0.12  | 66.38 <sup>b</sup> ± 26.55 | 47.71 <sup>e</sup> ± 10.06 | 76.03 <sup>b</sup> ± 13.64 | 35.48 <sup>b</sup> ± 17.29 | MR       |
| FB-26864             | 97.85 <sup>a</sup> ± 12.25 | 23.36 <sup>b</sup> ± 12.45 | 71.29 <sup>b</sup> ± 25.45 | 47.86 <sup>e</sup> ± 12.51 | 84.57 <sup>c</sup> ± 18.78 | 35.61 <sup>b</sup> ± 17.32 | MR       |
| FB-26399             | 87.00 <sup>b</sup> ± 14.25 | 23.42 <sup>b</sup> ± 0.25  | 78.82 <sup>f</sup> ± 25.02 | 48.06 <sup>e</sup> ± 22.99 | 82.91 <sup>c</sup> ± 5.78  | 35.74 <sup>b</sup> ± 17.42 | MR       |
| FB-28113             | 90.60 <sup>b</sup> ± 11.22 | 23.60 <sup>b</sup> ± 1.25  | 77.96 <sup>f</sup> ± 19.69 | 48.17 <sup>e</sup> ± 13.50 | 84.28 <sup>c</sup> ± 8.94  | 35.88 <sup>b</sup> ± 17.37 | MR       |
| FB-Ashebeke          | 98.50 <sup>a</sup> ± 10.25 | 23.58 <sup>b</sup> ± 0.08  | 65.41 <sup>b</sup> ± 20.27 | 48.18 <sup>e</sup> ± 21.14 | 81.96 <sup>c</sup> ± 23.40 | 35.88 <sup>b</sup> ± 17.39 | MR       |
| FB-28109             | 96.67 <sup>a</sup> ± 7.25  | 23.58 <sup>b</sup> ± 2.14  | 66.55 <sup>b</sup> ± 26.26 | 48.20 <sup>e</sup> ± 14.24 | 81.61 <sup>c</sup> ± 21.30 | 35.89 <sup>b</sup> ± 17.41 | MR       |
| FB-26877             | 96.67 <sup>a</sup> ± 8.22  | 25.08 <sup>b</sup> ± 1.45  | 68.06 <sup>b</sup> ± 31.31 | 48.69 <sup>e</sup> ± 17.07 | 82.37 <sup>c</sup> ± 20.23 | 36.89 <sup>b</sup> ± 16.69 | MR       |
| FB-28105             | 96.67 <sup>a</sup> ± 7.23  | 23.00 <sup>a</sup> ± 2.78  | 69.07 <sup>b</sup> ± 31.94 | 49.58 <sup>e</sup> ± 20.50 | 82.87 <sup>c</sup> ± 19.52 | 36.29 <sup>b</sup> ± 18.80 | MR       |
| FB-26876             | 95.00 <sup>a</sup> ± 9.36  | 22.83 <sup>a</sup> ± 0.47  | 65.61 <sup>b</sup> ± 23.05 | 50.67 <sup>e</sup> ± 4.23  | 80.30 <sup>c</sup> ± 20.78 | 36.75 <sup>b</sup> ± 19.69 | MR       |
| FB-212566            | 98.75 <sup>a</sup> ± 8.45  | 25.50 <sup>b</sup> ± 0.45  | 88.03 <sup>e</sup> ± 16.61 | 50.30 <sup>e</sup> ± 19.47 | 93.39 <sup>d</sup> ± 7.58  | 37.90 <sup>b</sup> ± 17.54 | MR       |



| Faba bean accessions | 2017 cropping season        |                            | 2018 cropping season       |                            | Combined mean              |                            | Reaction |
|----------------------|-----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------------------------|----------|
|                      | Incidence (%)<br>± SD       | Severity (%)<br>± SD       | Incidence (%) ±<br>SD      | Severity (%)<br>± SD       | Incidence (%)<br>± SD      | Severity (%)<br>± SD       |          |
| FB-26887             | 98.50 <sup>a</sup> ± 10.45  | 24.58 <sup>a</sup> ± 0.98  | 71.46 <sup>b</sup> ± 37.54 | 51.30 <sup>e</sup> ± 23.06 | 84.98 <sup>c</sup> ± 19.12 | 37.94 <sup>b</sup> ± 18.90 | MR       |
| FB-28103             | 96.33 <sup>a</sup> ± 16.56  | 22.17 <sup>a</sup> ± 6.07  | 86.04 <sup>e</sup> ± 22.64 | 55.62 <sup>f</sup> ± 27.66 | 91.19 <sup>d</sup> ± 7.28  | 38.90 <sup>b</sup> ± 23.66 | MR       |
| FB-25292             | 100.00 <sup>a</sup> ± 11.25 | 23.60 <sup>b</sup> ± 2.03  | 62.48 <sup>c</sup> ± 27.36 | 55.26 <sup>f</sup> ± 26.36 | 81.24 <sup>c</sup> ± 26.53 | 39.43 <sup>b</sup> ± 22.39 | MR       |
| FB-25022             | 96.00 <sup>a</sup> ± 15.47  | 23.60 <sup>b</sup> ± 5.04  | 87.70 <sup>e</sup> ± 18.87 | 55.30 <sup>f</sup> ± 13.63 | 91.85 <sup>d</sup> ± 5.87  | 39.45 <sup>b</sup> ± 22.42 | MR       |
| FB-Wayu              | 97.83 <sup>a</sup> ± 17.85  | 25.08 <sup>b</sup> ± 1.07  | 63.65 <sup>c</sup> ± 26.27 | 57.61 <sup>f</sup> ± 7.70  | 80.74 <sup>c</sup> ± 24.17 | 41.35 <sup>c</sup> ± 23.00 | MS       |
| FB-26881             | 96.67 <sup>a</sup> ± 19.52  | 24.67 <sup>b</sup> ± 2.09  | 60.90 <sup>c</sup> ± 22.20 | 57.80 <sup>f</sup> ± 4.52  | 78.78 <sup>b</sup> ± 25.30 | 41.23 <sup>c</sup> ± 23.42 | MS       |
| FB-28104             | 96.60 <sup>a</sup> ± 20.12  | 24.00 <sup>b</sup> ± 0.23  | 78.49 <sup>f</sup> ± 21.56 | 58.93 <sup>f</sup> ± 5.80  | 87.55 <sup>c</sup> ± 12.80 | 41.47 <sup>c</sup> ± 24.70 | MS       |
| Local check          | 98.19 <sup>a</sup> ± 32.56  | 42.43 <sup>c</sup> ± 12.54 | 71.42 <sup>b</sup> ± 27.79 | 44.05 <sup>d</sup> ± 20.43 | 84.81 <sup>c</sup> ± 18.93 | 43.50 <sup>c</sup> ± 13.88 | MS       |
| FB-26869             | 99.20 <sup>a</sup> ± 25.46  | 60.00 <sup>d</sup> ± 0.14  | 89.23 <sup>e</sup> ± 16.53 | 62.99 <sup>f</sup> ± 15.32 | 94.22 <sup>d</sup> ± 7.05  | 61.50 <sup>d</sup> ± 27.57 | S        |
| Mean                 | 94.94 <sup>a</sup> ± 12.35  | 22.74 <sup>a</sup> ± 2.05  | 66.89 <sup>b</sup> ± 29.53 | 38.73 <sup>d</sup> ± 12.72 | 80.80 <sup>c</sup> ± 19.68 | 30.61 <sup>b</sup> ± 12.86 | MR       |
| CV %                 | 12.00                       | 12.00                      | 44.15                      | 32.84                      | 24.36                      | 42.01                      |          |
| F-value              | 1.05                        | 7.56                       | 26.67                      | 38.34                      | 13.86                      | 22.95                      |          |
| LSD                  | 0.39                        | 0.00                       | 0.0003                     | 0.0002                     | 0.1951                     | 0.0001                     |          |

Means within a column followed by the same letter are not significantly different at  $p \geq 0.05$

CV: Coefficient of Variation, MR: Moderately Resistant, MS: Moderately Susceptible, S: Susceptible, SD: Standard Deviation

Table 3. Faba bean gall disease intensity on selected faba bean accessions during 2019 cropping season at different districts.

| Faba bean accessions | Angolelana Tera district        |                                   | Sululta district                  |                                   | Midakegn district                 |                                   |
|----------------------|---------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|-----------------------------------|
|                      | Incidence (%)                   | Severity (%)                      | Incidence (%)                     | Severity (%)                      | Incidence (%)                     | Severity (%)                      |
|                      | Mean $\pm$ SD                   | Mean $\pm$ SD                     | Mean $\pm$ SD                     | Mean $\pm$ SD                     | Mean $\pm$ SD                     | Mean $\pm$ SD                     |
| FB-Obse              | 40.42 <sup>c*</sup> $\pm$ 2.10  | 16.33 <sup>a*</sup> $\pm$ 13.28   | 31.02 <sup>b***</sup> $\pm$ 16.69 | 8.33 <sup>a***</sup> $\pm$ 3.33   | 55.33 <sup>a*</sup> $\pm$ 10.75   | 11.67 <sup>a*</sup> $\pm$ 2.89    |
| FB-28770             | 29.03 <sup>c*</sup> $\pm$ 12.25 | 16.56 <sup>a*</sup> $\pm$ 21.47   | 25.00 <sup>b*</sup> $\pm$ 3.30    | 11.67 <sup>a*</sup> $\pm$ 20.21   | 100.00 <sup>d***</sup> $\pm$ 0.00 | 45.00 <sup>e***</sup> $\pm$ 5.00  |
| FB-28107             | 48.61 <sup>a*</sup> $\pm$ 11.00 | 18.00 <sup>a*</sup> $\pm$ 19.40   | 9.78 <sup>a***</sup> $\pm$ 5.98   | 12.89 <sup>a*</sup> $\pm$ 6.71    | 98.33 <sup>d***</sup> $\pm$ 31.79 | 36.67 <sup>d**</sup> $\pm$ 17.89  |
| FB-Hachalu           | 41.30 <sup>a*</sup> $\pm$ 12.30 | 21.67 <sup>b*</sup> $\pm$ 10.69   | 23.17 <sup>b**</sup> $\pm$ 2.45   | 6.67 <sup>a***</sup> $\pm$ 7.66   | 74.89 <sup>b***</sup> $\pm$ 9.11  | 20.00 <sup>b*</sup> $\pm$ 18.03   |
| FB-26875             | 44.72 <sup>a*</sup> $\pm$ 3.12  | 28.33 <sup>b*</sup> $\pm$ 4.71    | 32.09 <sup>b*</sup> $\pm$ 30.86   | 12.78 <sup>a***</sup> $\pm$ 14.96 | 76.67 <sup>b**</sup> $\pm$ 20.55  | 33.33 <sup>c*</sup> $\pm$ 26.25   |
| FB-26873             | 50.61 <sup>a*</sup> $\pm$ 1.03  | 30.00 <sup>c*</sup> $\pm$ 7.64    | 42.96 <sup>c*</sup> $\pm$ 6.80    | 17.22 <sup>b***</sup> $\pm$ 10.18 | 88.56 <sup>c**</sup> $\pm$ 19.82  | 28.33 <sup>b*</sup> $\pm$ 20.21   |
| FB-26883             | 69.08 <sup>b*</sup> $\pm$ 15.23 | 31.83 <sup>c*</sup> $\pm$ 17.48   | 31.92 <sup>b***</sup> $\pm$ 16.56 | 15.55 <sup>b***</sup> $\pm$ 12.52 | 80.96 <sup>c*</sup> $\pm$ 22.90   | 40.00 <sup>d***</sup> $\pm$ 17.32 |
| FB-Numan             | 33.33 <sup>c*</sup> $\pm$ 18.02 | 31.00 <sup>c*</sup> $\pm$ 15.25   | 22.50 <sup>b*</sup> $\pm$ 3.39    | 12.22 <sup>a***</sup> $\pm$ 11.10 | 88.67 <sup>c**</sup> $\pm$ 19.63  | 39.00 <sup>d*</sup> $\pm$ 20.00   |
| FB-26867             | 33.94 <sup>c*</sup> $\pm$ 1.23  | 34.17 <sup>c**</sup> $\pm$ 11.25  | 17.41 <sup>a***</sup> $\pm$ 5.95  | 11.11 <sup>a*</sup> $\pm$ 16.36   | 77.89 <sup>b***</sup> $\pm$ 38.30 | 15.00 <sup>a*</sup> $\pm$ 8.66    |
| FB-Tumsa             | 49.62 <sup>a*</sup> $\pm$ 9.00  | 36.00 <sup>c***</sup> $\pm$ 29.18 | 35.19 <sup>b***</sup> $\pm$ 3.64  | 18.33 <sup>b*</sup> $\pm$ 16.10   | 73.33 <sup>b***</sup> $\pm$ 23.09 | 36.67 <sup>d**</sup> $\pm$ 11.55  |
| FB-26885             | 43.18 <sup>a*</sup> $\pm$ 2.25  | 42.50 <sup>d*</sup> $\pm$ 3.89    | 36.11 <sup>b*</sup> $\pm$ 29.29   | 22.22 <sup>b***</sup> $\pm$ 28.70 | 100.00 <sup>d***</sup> $\pm$ 0.00 | 46.67 <sup>e*</sup> $\pm$ 15.28   |
| FB-25280             | 74.44 <sup>b*</sup> $\pm$ 14.02 | 43.10 <sup>d*</sup> $\pm$ 6.11    | 66.67 <sup>c*</sup> $\pm$ 0.00    | 12.22 <sup>a***</sup> $\pm$ 14.58 | 93.33 <sup>c**</sup> $\pm$ 11.55  | 30.00 <sup>c***</sup> $\pm$ 26.46 |
| FB-26872             | 52.22 <sup>a*</sup> $\pm$ 1.23  | 46.33 <sup>d*</sup> $\pm$ 32.72   | 14.81 <sup>a***</sup> $\pm$ 19.60 | 6.33 <sup>a***</sup> $\pm$ 7.77   | 85.57 <sup>c***</sup> $\pm$ 17.09 | 23.33 <sup>b***</sup> $\pm$ 11.55 |
| FB- Coll-0038        | 35.56 <sup>c*</sup> $\pm$ 4.58  | 48.89 <sup>d*</sup> $\pm$ 27.10   | 29.17 <sup>b*</sup> $\pm$ 5.89    | 6.67 <sup>a***</sup> $\pm$ 6.02   | 81.56 <sup>c**</sup> $\pm$ 31.94  | 40.00 <sup>d*</sup> $\pm$ 20.00   |
| Local check          | 43.33 <sup>a*</sup> $\pm$ 10.25 | 63.00 <sup>e*</sup> $\pm$ 1.89    | 25.59 <sup>b***</sup> $\pm$ 5.59  | 22.38 <sup>b***</sup> $\pm$ 22.38 | 82.78 <sup>c***</sup> $\pm$ 20.57 | 31.67 <sup>c*</sup> $\pm$ 14.43   |
| Average              | 45.96 <sup>a*</sup> $\pm$ 12.52 | 38.76 <sup>c*</sup> $\pm$ 24.41   | 29.56 <sup>b***</sup> $\pm$ 13.44 | 13.11 <sup>a***</sup> $\pm$ 5.20  | 83.86 <sup>c***</sup> $\pm$ 11.87 | 31.89 <sup>c*</sup> $\pm$ 58.80   |

Means within a column followed by the same letter are not significantly different at  $p \geq 0.05$ , Means across the district within a row followed by the same number of \* are not significantly different at  $p \geq 0.05$ .

### **Combined mean analysis of gall disease intensity over the three cropping seasons**

The interaction effect of faba bean accessions with districts and cropping season showed significant difference ( $p \leq 0.05$ ) in gall disease incidence and severity. In combined mean analysis over year (2017–2019), the DSI of 14 accessions and local check was recorded between 16.50–41.8% (Table 4).

The lowest DSI was recorded on FB-Obse (DSI = 16.50%) followed by FB-Hachalu (DSI = 17.83%), FB-26872 (DSI = 18.84%), FB-26873 (DSI = 19.61%), and FB-28107 (DSI = 19.85%) over the three districts. The local check exhibited higher DSI (41.8%) than the remaining accessions (Table 4). For the reaction of the 14 accessions classified as moderately resistant, there was significant variation on their reaction level against gall disease. The combined mean analysis of DSI of gall disease over the faba bean accessions revealed that the average maximum DSI (26.3%) was recorded during 2019 cropping seasons followed by 2017 (DSI = 21.8%) and 2018 (DSI = 20.3%) (Table 4).

### **Combined gall disease severity over districts**

Over the three cropping seasons, the highest DSI (31.58%) was recorded at Angolelana Tera followed by Midakegn (25.53%), and Sululta (13.29%) districts (Fig. 1). The lowest DSI was recorded on FB-Hachalu (9.51%), FB-Obse (9.72%), FB-26872 (9.77%), and FB-26873 (10.50%) at Sululta, whereas the lowest DSI was recorded on FB-Obse (15.20%), FB-Hachalu (18.55%), FB-26872 (20.22%), and FB-Tumsa (21.08%) at Midakegn district. Similarly, the lowest DSI was recorded on FB-Obse (18.17%), FB-26885 (24.58%), FB-Tumsa (25.46%), and FB-Numan (25.50%) at Angolelana Tera district (Fig. 1).

Table 4. Average gall disease intensity of selected faba bean accessions over experimental sites during three cropping seasons.

|                      | 2017 cropping season     |                          | 2018 cropping season     |                          | 2019 cropping season     |                          | Combined mean            |                         |          |
|----------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|-------------------------|----------|
| Faba bean accessions | Incidence (%) ± SD       | Severity (%) ± SD        | Incidence (%) ± SD       | Severity (%) ± SD        | Incidence (%) ± SD       | Severity (%) ± SD        | Incidence (%) ± SD       | Severity (%) ± SD       | Reaction |
| FB-Obse              | 98.6 <sup>a</sup> ± 11.2 | 21.1 <sup>a</sup> ± 1.2  | 53.3 <sup>c</sup> ± 39.2 | 16.3 <sup>b</sup> ± 3.7  | 42.3 <sup>e</sup> ± 12.3 | 12.1 <sup>f</sup> ± 4.0  | 64.7 <sup>g</sup> ± 29.9 | 16.5 <sup>b</sup> ± 4.5 | MR       |
| FB-Hachalu           | 97.7 <sup>a</sup> ± 10.5 | 21.2 <sup>a</sup> ± 0.5  | 51.9 <sup>c</sup> ± 44.9 | 16.2 <sup>b</sup> ± 3.6  | 46.5 <sup>e</sup> ± 26.2 | 16.1 <sup>b</sup> ± 8.2  | 65.3 <sup>g</sup> ± 28.1 | 17.8 <sup>b</sup> ± 2.9 | MR       |
| FB-26872             | 88.8 <sup>b</sup> ± 3.0  | 16.8 <sup>b</sup> ± 0.4  | 50.8 <sup>c</sup> ± 42.1 | 14.4 <sup>b</sup> ± 4.9  | 50.9 <sup>e</sup> ± 35.4 | 25.3 <sup>g</sup> ± 20.1 | 63.5 <sup>g</sup> ± 21.9 | 18.8 <sup>b</sup> ± 5.8 | MR       |
| FB-26873             | 91.2 <sup>b</sup> ± 8.3  | 17.0 <sup>b</sup> ± 5.1  | 58.3 <sup>d</sup> ± 36.7 | 16.6 <sup>b</sup> ± 4.8  | 60.7 <sup>d</sup> ± 24.4 | 25.2 <sup>g</sup> ± 6.9  | 70.1 <sup>e</sup> ± 18.3 | 19.6 <sup>b</sup> ± 4.8 | MR       |
| FB-28107             | 97.2 <sup>a</sup> ± 12.1 | 21.1 <sup>a</sup> ± 1.3  | 60.3 <sup>d</sup> ± 34.1 | 15.9 <sup>b</sup> ± 3.8  | 52.2 <sup>c</sup> ± 44.4 | 22.5 <sup>a</sup> ± 12.5 | 69.9 <sup>e</sup> ± 23.9 | 19.9 <sup>b</sup> ± 3.5 | MR       |
| FB-Tumsa             | 80.0 <sup>d</sup> ± 11.3 | 20.2 <sup>a</sup> ± 1.6  | 52.5 <sup>c</sup> ± 30.7 | 13.2 <sup>b</sup> ± 1.0  | 52.7 <sup>c</sup> ± 18.0 | 30.3 <sup>d</sup> ± 1.4  | 61.7 <sup>g</sup> ± 5.8  | 21.2 <sup>a</sup> ± 2.6 | MR       |
| FB-26867             | 91.0 <sup>b</sup> ± 4.0  | 23.7 <sup>a</sup> ± 1.0  | 46.0 <sup>a</sup> ± 25.8 | 21.1 <sup>a</sup> ± 1.6  | 43.1 <sup>a</sup> ± 21.3 | 20.1 <sup>a</sup> ± 2.3  | 60.0 <sup>g</sup> ± 6.9  | 21.6 <sup>a</sup> ± 0.8 | MR       |
| FB-Numan             | 90.1 <sup>b</sup> ± 1.3  | 21.5 <sup>a</sup> ± 1.5  | 59.5 <sup>d</sup> ± 23.9 | 16.5 <sup>b</sup> ± 2.9  | 48.2 <sup>b</sup> ± 25.5 | 27.4 <sup>g</sup> ± 3.7  | 65.9 <sup>g</sup> ± 1.7  | 21.8 <sup>a</sup> ± 2.5 | MR       |
| FB-28770             | 85.0 <sup>c</sup> ± 1.3  | 22.4 <sup>a</sup> ± 1.6  | 56.3 <sup>d</sup> ± 17.4 | 18.4 <sup>c</sup> ± 2.0  | 51.3 <sup>c</sup> ± 12.2 | 24.4 <sup>g</sup> ± 7.9  | 64.2 <sup>g</sup> ± 8.2  | 21.8 <sup>a</sup> ± 1.0 | MR       |
| FB-26875             | 95.2 <sup>a</sup> ± 7.2  | 21.6 <sup>a</sup> ± 1.8  | 64.7 <sup>e</sup> ± 11.2 | 21.1 <sup>a</sup> ± 6.1  | 51.2 <sup>c</sup> ± 12.9 | 24.8 <sup>g</sup> ± 1.7  | 70.3 <sup>e</sup> ± 2.6  | 22.5 <sup>a</sup> ± 1.0 | MR       |
| FB-26883             | 91.1 <sup>b</sup> ± 2.3  | 18.2 <sup>b</sup> ± 1.6  | 60.3 <sup>d</sup> ± 12.8 | 20.9 <sup>a</sup> ± 1.8  | 60.7 <sup>d</sup> ± 5.6  | 29.1 <sup>d</sup> ± 2.4  | 70.7 <sup>e</sup> ± 7.7  | 22.7 <sup>a</sup> ± 3.7 | MR       |
| FB-COLL-0038         | 96.8 <sup>a</sup> ± 8.5  | 21.1 <sup>a</sup> ± 0.3  | 71.6 <sup>f</sup> ± 7.1  | 21.0 <sup>a</sup> ± 5.0  | 48.8 <sup>b</sup> ± 8.6  | 31.9 <sup>d</sup> ± 2.3  | 72.4 <sup>e</sup> ± 4.0  | 24.6 <sup>g</sup> ± 2.3 | MR       |
| FB-25280             | 95.0 <sup>a</sup> ± 1.9  | 17.8 <sup>b</sup> ± 1.5  | 53.7 <sup>c</sup> ± 23.5 | 28.2 <sup>d</sup> ± 7.7  | 78.6 <sup>e</sup> ± 3.7  | 28.4 <sup>d</sup> ± 5.5  | 75.6 <sup>e</sup> ± 2.8  | 24.8 <sup>g</sup> ± 4.1 | MR       |
| FB-26885             | 98.8 <sup>a</sup> ± 12.5 | 21.8 <sup>a</sup> ± 4.1  | 57.4 <sup>d</sup> ± 36.6 | 20.9 <sup>a</sup> ± 6.7  | 59.8 <sup>d</sup> ± 35.0 | 37.1 <sup>e</sup> ± 13.1 | 72.0 <sup>e</sup> ± 23.3 | 26.6 <sup>g</sup> ± 9.1 | MR       |
| Local check          | 98.2 <sup>a</sup> ± 32.6 | 42.4 <sup>e</sup> ± 12.5 | 71.4 <sup>f</sup> ± 27.8 | 44.1 <sup>e</sup> ± 20.4 | 50.6 <sup>b</sup> ± 29.3 | 39.0 <sup>e</sup> ± 21.3 | 73.4 <sup>e</sup> ± 23.9 | 41.8 <sup>e</sup> ± 2.6 | MS       |
| Average              | 92.9 <sup>b</sup> ± 5.5  | 21.8 <sup>a</sup> ± 6.0  | 57.9 <sup>d</sup> ± 7.2  | 20.3 <sup>a</sup> ± 7.6  | 53.1 <sup>b</sup> ± 8.9  | 26.3 <sup>g</sup> ± 7.1  | 67.9 <sup>e</sup> ± 4.7  | 22.8 <sup>a</sup> ± 5.8 | MR       |
| CV (%)               | 5.92                     | 27.62                    | 12.46                    | 37.19                    | 16.77                    | 27.04                    | 6.84                     | 25.91                   | MR       |
| LSD                  | 0.39                     | 0                        | 0.0003                   | 0.0002                   | 0.0008                   | 0.0006                   | 0.007                    | 0.0025                  | MR       |

The same letter in the same column indicates absence of significant difference ( $p > 0.05$ ), SD = standard deviations, CV = Coefficient of Variation, LSD = Least Significant difference, MR = Moderately resistant, MS = Moderately susceptible




Fig.1. Gall disease severity on faba bean accessions at various localities.

### Area Under Disease Progress Curve (AUDPC) and apparent infection rate

The date of gall disease incidence on faba bean accessions varied across the districts over the three cropping seasons (2017–2019) (Fig. 2). At Angolelana Tera, the gall disease incidence was observed between 15 and 38 days, whereas at Sululta district it was observed between 38 and 77 days after crop emergence. Higher DSI was recorded at Angolelana Tera (DSI = 13.68–47.29%) than the Sululta district throughout the cropping seasons. The fastest DSI rate (74.88–100%) was recorded between 38–77 days after emergence at both districts. At Angolelana Tera, and Sululta districts, a slow DSI rate (55.78–49.78%) was observed starting from 77 days, and become constant between 103–115 days after the crop emergence (Fig. 2). Similarly, higher AUDPC was recorded at Angolelana Tera (1616.97) than Sululta (1238.46) district with maximum infection rate ( $r = 0.02$ ) and coefficient of determination ( $R^2 = 88\%$ ) (Table 5).




Fig. 2. Gall disease progress curve at Angolelana Tera and Sululta districts. Vertical line on the graphs indicates standard error.

## Analysis of agronomic parameters versus gall disease intensity

### Seed germination percentage of faba bean accessions over study sites

The maximum seed emergence percentage was recorded on FB-Obse at Sululta (98.26%) district. FB-Coll-0038 accession showed significantly higher field emergence efficiency than others at Midakegn (96.30%) district (Table 6). The percentage germination of seeds in the field at Angolelana Tera was significantly lower than at Sululta and Midakegn districts. There was no significant difference ( $p \geq 0.05$ ) between Sululta and Midakegn districts in a combined mean average percentage germination of faba bean seeds. The highest combined faba bean accession mean average germination percentage was recorded at Sululta (90.99%) followed by Midakegn (85.00%), and Angolelana Tera (47.73%). The combined mean average germination analysis of faba bean accessions over the three districts revealed that the highest seed germination percentage was recorded on FB-26873 (79.85%), followed by FB-Obse (79.30%), and FB-Hachalu (79.09%) (Table 6).

Table 5. AUDPC and apparent infection rate of faba bean gall disease over three cropping seasons (2017–2019).

| Faba bean accessions | Angolelana Tera district      |                              |                                                    | Sululta district                |                             |                                                    |
|----------------------|-------------------------------|------------------------------|----------------------------------------------------|---------------------------------|-----------------------------|----------------------------------------------------|
|                      | Mean AUDPC                    | Apparent infection rate (r)  | Coefficient of determination (R <sup>2</sup> ) (%) | Mean AUDPC                      | Apparent infection rate (r) | Coefficient of determination (R <sup>2</sup> ) (%) |
|                      | Mean ± SD                     | Mean ± SD                    |                                                    | Mean ± SD                       | Mean ± SD                   |                                                    |
| FB-Obse              | 815.50 <sup>a*</sup> ± 68.68  | 0.01 <sup>a*</sup> ± 0.1     | 92                                                 | 523.89 <sup>a***</sup> ± 43.08  | 0.01 <sup>b*</sup> ± 0.6    | 83                                                 |
| FB-28770             | 763.50 <sup>c*</sup> ± 31.09  | 0.02 <sup>a*</sup> ± 0.5     | 82                                                 | 1843.03 <sup>d***</sup> ± 13.05 | 0.01 <sup>b***</sup> ± 0.3  | 80                                                 |
| FB-28107             | 815.51 <sup>c*</sup> ± 43.90  | 0.01 <sup>a*</sup> ± 0.04    | 86                                                 | 831.86 <sup>c*</sup> ± 49.04    | 0.01 <sup>b*</sup> ± 0.07   | 85                                                 |
| FB-Hachalu           | 1072.60 <sup>a*</sup> ± 54.08 | 0.01 <sup>a*</sup> ± 0.1     | 91                                                 | 140.02 <sup>b***</sup> ± 50.07  | 0.01 <sup>b*</sup> ± 0.7    | 89                                                 |
| FB-26875             | 1373.25 <sup>a*</sup> ± 42.01 | 0.02 <sup>a*</sup> ± 0.1     | 89                                                 | 1015.36 <sup>c***</sup> ± 13.03 | 0.01 <sup>b*</sup> ± 0.8    | 81                                                 |
| FB-26873             | 1417.50 <sup>a*</sup> ± 10.04 | 0.01 <sup>a*</sup> ± 0.01    | 93                                                 | 910.00 <sup>c***</sup> ± 63.99  | 0.003 <sup>d*</sup> ± 0.01  | 81                                                 |
| FB-26883             | 1581.01 <sup>a*</sup> ± 9.20  | 0.02 <sup>a*</sup> ± 0.2     | 96                                                 | 1115.10 <sup>c***</sup> ± 98.00 | 0.06 <sup>c***</sup> ± 0.2  | 86                                                 |
| FB-Numan             | 1620.75 <sup>a*</sup> ± 92.10 | 0.002 <sup>b***</sup> ± 0.10 | 48                                                 | 1534.52 <sup>d*</sup> ± 42.03   | 0.02 <sup>a*</sup> ± 0.10   | 86                                                 |
| FB-26867             | 2128.75 <sup>b*</sup> ± 95.05 | 0.02 <sup>a*</sup> ± 0.01    | 94                                                 | 548.03 <sup>a***</sup> ± 92.00  | 0.01 <sup>b*</sup> ± 0.12   | 83                                                 |
| FB-Tumsa             | 1746.00 <sup>a*</sup> ± 63.06 | 0.01 <sup>a*</sup> ± 0.11    | 92                                                 | 1044.16 <sup>c***</sup> ± 51.20 | 0.01 <sup>b***</sup> ± 0.11 | 85                                                 |
| FB-26885             | 2004.05 <sup>b*</sup> ± 19.03 | 0.02 <sup>a*</sup> ± 0.00    | 86                                                 | 2135.10 <sup>d*</sup> ± 69.02   | 0.01 <sup>b***</sup> ± 0.01 | 81                                                 |
| FB-25280             | 2114.50 <sup>b*</sup> ± 13.57 | 0.02 <sup>a*</sup> ± 0.01    | 92                                                 | 974.69 <sup>c***</sup> ± 87.02  | 0.01 <sup>b***</sup> ± 0.00 | 80                                                 |
| FB-26872             | 2173.00 <sup>b*</sup> ± 8.03  | 0.08 <sup>a*</sup> ± 0.10    | 86                                                 | 682.50 <sup>a***</sup> ± 80.01  | 0.01 <sup>b*</sup> ± 0.11   | 80                                                 |
| FB- Coll-0038        | 2322.50 <sup>b*</sup> ± 81.03 | 0.04 <sup>a*</sup> ± 0.14    | 88                                                 | 4262.50 <sup>d***</sup> ± 80.04 | 0.01 <sup>b***</sup> ± 0.21 | 82                                                 |
| Local check          | 2961.75 <sup>b*</sup> ± 94.41 | 0.02 <sup>a*</sup> ± 0.01    | 90                                                 | 1015.36 <sup>c***</sup> ± 22.06 | 0.01 <sup>b*</sup> ± 0.18   | 80                                                 |
| Average              | 1616.97 <sup>a*</sup> ± 67.03 | 0.02 <sup>a*</sup> ± 0.11    | 88                                                 | 1238.46 <sup>c***</sup> ± 78.05 | 0.01 <sup>b***</sup> ± 0.13 | 79                                                 |

Similar letters within the same column indicate an absence of significant difference ( $P > 0.05$ ), means across the district within a row followed by the same number of \* are not significantly different at  $p > 0.05$ , SD = standard deviation

Table 6. Seed germination of faba bean accessions at different districts.

| Faba bean accessions | Angolelana Tera district       | Sululta district                  | Midakegn district                  | Combined                          |
|----------------------|--------------------------------|-----------------------------------|------------------------------------|-----------------------------------|
|                      | Germination (%)                | Germination (%)                   | Germination (%)                    | Germination (%)                   |
|                      | Mean $\pm$ SD                  | Mean $\pm$ SD                     | Mean $\pm$ SD                      | Mean $\pm$ SD                     |
| FB-Obse              | 61.85 <sup>a*</sup> $\pm$ 2.08 | 98.26 <sup>a****</sup> $\pm$ 3.06 | 77.78 <sup>c***</sup> $\pm$ 22.22  | 79.30 <sup>a**</sup> $\pm$ 18.25  |
| FB-28770             | 57.00 <sup>a*</sup> $\pm$ 1.73 | 77.04 <sup>c**</sup> $\pm$ 2.08   | 85.19 <sup>b***</sup> $\pm$ 12.83  | 73.08 <sup>ab**</sup> $\pm$ 14.51 |
| FB-28107             | 44.44 <sup>b*</sup> $\pm$ 3.61 | 90.56 <sup>b****</sup> $\pm$ 1.00 | 85.19 <sup>b****</sup> $\pm$ 30.40 | 73.40 <sup>ab**</sup> $\pm$ 25.22 |
| FB-Hachalu           | 58.00 <sup>a*</sup> $\pm$ 2.08 | 94.07 <sup>a****</sup> $\pm$ 1.53 | 85.19 <sup>b**</sup> $\pm$ 6.42    | 79.09 <sup>a**</sup> $\pm$ 18.79  |
| FB-26875             | 48.00 <sup>b*</sup> $\pm$ 0.00 | 94.07 <sup>a****</sup> $\pm$ 4.04 | 81.48 <sup>c**</sup> $\pm$ 13.86   | 74.52 <sup>ab**</sup> $\pm$ 23.81 |
| FB-26873             | 55.00 <sup>a*</sup> $\pm$ 1.53 | 91.96 <sup>b****</sup> $\pm$ 0.58 | 92.59 <sup>a****</sup> $\pm$ 6.42  | 79.85 <sup>a**</sup> $\pm$ 21.52  |
| FB-26883             | 48.15 <sup>b*</sup> $\pm$ 0.71 | 94.78 <sup>a****</sup> $\pm$ 1.00 | 81.48 <sup>c**</sup> $\pm$ 6.42    | 74.80 <sup>ab**</sup> $\pm$ 24.02 |
| FB-Numan             | 50.00 <sup>b*</sup> $\pm$ 2.12 | 97.67 <sup>a****</sup> $\pm$ 1.73 | 77.78 <sup>c**</sup> $\pm$ 29.40   | 75.15 <sup>ab**</sup> $\pm$ 23.94 |
| FB-26867             | 48.15 <sup>b*</sup> $\pm$ 0.71 | 79.26 <sup>c**</sup> $\pm$ 3.51   | 88.89 <sup>b**</sup> $\pm$ 11.11   | 72.10 <sup>b**</sup> $\pm$ 21.29  |
| FB-Tumsa             | 52.00 <sup>b*</sup> $\pm$ 1.73 | 97.67 <sup>a****</sup> $\pm$ 1.00 | 81.48 <sup>c**</sup> $\pm$ 12.83   | 77.05 <sup>a**</sup> $\pm$ 23.16  |
| FB-26885             | 37.04 <sup>c*</sup> $\pm$ 4.24 | 93.85 <sup>b****</sup> $\pm$ 0.58 | 88.89 <sup>b****</sup> $\pm$ 11.11 | 73.26 <sup>ab**</sup> $\pm$ 31.47 |
| FB-25280             | 38.00 <sup>c*</sup> $\pm$ 1.73 | 87.04 <sup>b****</sup> $\pm$ 2.89 | 81.48 <sup>c****</sup> $\pm$ 6.42  | 68.84 <sup>c**</sup> $\pm$ 26.85  |
| FB-26872             | 45.00 <sup>b*</sup> $\pm$ 0.58 | 96.1b <sup>a****</sup> $\pm$ 2.31 | 88.89 <sup>b****</sup> $\pm$ 11.11 | 76.66 <sup>a**</sup> $\pm$ 27.66  |
| FB- Coll-0038        | 38.00 <sup>c*</sup> $\pm$ 1.73 | 87.04 <sup>b****</sup> $\pm$ 3.06 | 96.30 <sup>a****</sup> $\pm$ 6.42  | 73.78 <sup>ab**</sup> $\pm$ 31.33 |
| Local check          | 35.00 <sup>c*</sup> $\pm$ 1.41 | 85.37 <sup>b****</sup> $\pm$ 1.53 | 88.89 <sup>b****</sup> $\pm$ 11.11 | 69.75 <sup>b**</sup> $\pm$ 30.15  |
| Average              | 47.73 <sup>b*</sup> $\pm$ 1.02 | 90.99 <sup>a****</sup> $\pm$ 2.14 | 85.00 <sup>b****</sup> $\pm$ 0.89  | 74.71 <sup>ab**</sup> $\pm$ 23.44 |

Means within a column followed by the same letter are not significantly different at  $p \geq 0.05$ , Means across the district within a row followed by the same number of \* are not significantly different at  $p \geq 0.05$



## **Yield parameters**

### **Analysis of yield parameters in 2018 cropping season**

Faba bean gall disease significantly affected ( $p < 0.05$ ) yield parameters in various experimental sites. In 2018 cropping season, the highest number of pods per plant was obtained from FB-25280 (6.07 pods/plant) followed by FB-28770 (5.73 pods/plant) at Sululta and Angolelana Tera districts, respectively (Table 7). FB-26867 accession had produced maximum number of seeds per pod (3.00 seeds/pod) at Sululta districts. FB-26875 accession had exhibited highest number of seeds per pod (1.98 seeds/pod) among others at Angolelana Tera district. On another hand, FB-26872 produced maximum 100 seeds weight at Midakegn (97.97 g) and Sululta (90.87 g) districts. But, maximum 100 seeds weight was recorded on FB-Numan (72.53 g) at Angolelana Tera district. In a combined analysis over the faba bean accessions, the highest 100 seed weight was recorded at Midakegn (77.67 g) followed by Sululta (71.37 g) and Angolelana Tera (19.98 g) districts (Table 7).

### **Analysis of yield parameters in 2019 cropping season**

The combined mean analysis of faba bean accessions at each experimental site revealed that significantly ( $p < 0.05$ ) higher pods per plant was recorded at Sululta (9.26 pods/plant) than Angolelana Tera (1.55 pods/plant) district. Similar patterns were followed in number of pods per plant and number of seeds per pod (Table 8). The highest number of pods per plant was recorded on FB-Numan both at Sululta (14.18 pods/plant) and Angolelana Tera (5.31 pods/plant) districts. The number of seeds per pod was not significantly different ( $p > 0.05$ ) among the tested accessions at Sululta district, while the maximum seeds per pod recorded on FB-26875 accession (11.02 seeds/pod) at Angolelana Tera district. The maximum 100 seeds weight was recorded on FB-Obse (78.69 g) at Sululta followed by FB-Hachalu (75 g), and FB-Tumsa (55.34 g) at Midakegn and Angolelana Tera districts, respectively (Table 8).

Table 7. Average yield parameters of some faba bean accessions at different experimental sites during 2018 cropping season.

| Faba bean accessions | Field experimental sites     |                              |                                |                              |                              |                                |                                |
|----------------------|------------------------------|------------------------------|--------------------------------|------------------------------|------------------------------|--------------------------------|--------------------------------|
|                      | Angolelana Tera              |                              |                                | Sululta                      |                              |                                | Midakegn                       |
|                      | Pod/ plant                   | Seed/pod                     | 100 seed weight (g)            | Pod/ plant                   | Seed/pod                     | 100 seed weight (g)            | 100 seed weight (g)            |
|                      | Mean $\pm$ SD                | Mean $\pm$ SD                | Mean $\pm$ SD                  | Mean $\pm$ SD                | Mean $\pm$ SD                | Mean $\pm$ SD                  | Mean $\pm$ SD                  |
| FB-Obse              | 2.12 <sup>c</sup> $\pm$ 0.12 | 0.30 <sup>b</sup> $\pm$ 0.01 | 54.14 <sup>b</sup> $\pm$ 11.36 | 1.85 <sup>e</sup> $\pm$ 0.12 | 2.00 <sup>b</sup> $\pm$ 0.69 | 86.31 <sup>a</sup> $\pm$ 10.25 | 90.81 <sup>a</sup> $\pm$ 10.23 |
| FB-28770             | 5.73 <sup>a</sup> $\pm$ 0.21 | 1.72 <sup>a</sup> $\pm$ 0.23 | 20.44 <sup>d</sup> $\pm$ 8.06  | 3.00 <sup>c</sup> $\pm$ 0.23 | 1.63 <sup>b</sup> $\pm$ 0.78 | 65.33 <sup>c</sup> $\pm$ 11.36 | 70.06 <sup>c</sup> $\pm$ 12.95 |
| FB-28107             | 1.89 <sup>c</sup> $\pm$ 0.54 | 0.74 <sup>b</sup> $\pm$ 0.21 | 34.44 <sup>c</sup> $\pm$ 8.34  | 2.63 <sup>d</sup> $\pm$ 0.45 | 2.11 <sup>b</sup> $\pm$ 0.36 | 69.22 <sup>c</sup> $\pm$ 15.67 | 82.59 <sup>b</sup> $\pm$ 12.65 |
| FB-Hachalu           | 3.82 <sup>b</sup> $\pm$ 1.20 | 1.80 <sup>a</sup> $\pm$ 0.43 | 36.60 <sup>c</sup> $\pm$ 9.36  | 1.80 <sup>e</sup> $\pm$ 0.69 | 2.73 <sup>a</sup> $\pm$ 0.56 | 71.25 <sup>c</sup> $\pm$ 13.96 | 84.31 <sup>b</sup> $\pm$ 15.60 |
| FB-26875             | 4.07 <sup>b</sup> $\pm$ 2.00 | 1.98 <sup>a</sup> $\pm$ 0.56 | 20.68 <sup>d</sup> $\pm$ 5.89  | 4.20 <sup>b</sup> $\pm$ 0.47 | 1.41 <sup>c</sup> $\pm$ 0.45 | 82.73 <sup>b</sup> $\pm$ 16.58 | 80.73 <sup>b</sup> $\pm$ 8.96  |
| FB-26873             | 1.76 <sup>c</sup> $\pm$ 0.01 | 1.34 <sup>a</sup> $\pm$ 0.53 | 14.39 <sup>e</sup> $\pm$ 4.23  | 3.02 <sup>c</sup> $\pm$ 0.25 | 1.77 <sup>b</sup> $\pm$ 0.98 | 80.13 <sup>b</sup> $\pm$ 17.6  | 79.04 <sup>b</sup> $\pm$ 11.84 |
| FB-26883             | 5.59 <sup>a</sup> $\pm$ 1.30 | 0.68 <sup>b</sup> $\pm$ 0.12 | 17.67 <sup>e</sup> $\pm$ 3.78  | 3.64 <sup>a</sup> $\pm$ 0.23 | 1.66 <sup>b</sup> $\pm$ 0.74 | 66.95 <sup>c</sup> $\pm$ 5.98  | 72.80 <sup>c</sup> $\pm$ 7.98  |
| FB-Numan             | 2.69 <sup>c</sup> $\pm$ 0.23 | 1.60 <sup>a</sup> $\pm$ 0.32 | 72.53 <sup>a</sup> $\pm$ 20.7  | 2.38 <sup>d</sup> $\pm$ 0.15 | 2.53 <sup>a</sup> $\pm$ 0.92 | 65.64 <sup>c</sup> $\pm$ 7.95  | 71.87 <sup>c</sup> $\pm$ 8.96  |
| FB-26867             | 1.68 <sup>c</sup> $\pm$ 0.41 | 0.26 <sup>b</sup> $\pm$ 0.01 | 15.50 <sup>e</sup> $\pm$ 2.89  | 2.13 <sup>d</sup> $\pm$ 0.45 | 3.00 <sup>a</sup> $\pm$ 0.85 | 79.61 <sup>b</sup> $\pm$ 12.30 | 71.73 <sup>c</sup> $\pm$ 8.65  |
| FB-Tumsa             | 4.14 <sup>b</sup> $\pm$ 1.52 | 1.53 <sup>a</sup> $\pm$ 0.23 | 36.18 <sup>c</sup> $\pm$ 21.36 | 1.70 <sup>e</sup> $\pm$ 0.18 | 2.77 <sup>a</sup> $\pm$ 0.45 | 70.00 <sup>c</sup> $\pm$ 12.65 | 88.85 <sup>a</sup> $\pm$ 4.68  |
| FB-26885             | 2.27 <sup>c</sup> $\pm$ 0.21 | 1.89 <sup>a</sup> $\pm$ 0.25 | 64.90 <sup>a</sup> $\pm$ 11.58 | 3.28 <sup>c</sup> $\pm$ 0.14 | 1.90 <sup>b</sup> $\pm$ 0.35 | 56.16 <sup>d</sup> $\pm$ 14.60 | 73.12 <sup>c</sup> $\pm$ 5.67  |
| FB-25280             | 0.00 <sup>d</sup> $\pm$ 0.00 | 0.00 <sup>c</sup> $\pm$ 0.00 | 0.00 <sup>f</sup> $\pm$ 0.00   | 6.07 <sup>a</sup> $\pm$ 0.16 | 1.91 <sup>b</sup> $\pm$ 0.36 | 70.27 <sup>c</sup> $\pm$ 10.90 | 72.88 <sup>c</sup> $\pm$ 10.95 |
| FB-26872             | 3.95 <sup>b</sup> $\pm$ 1.23 | 1.67 <sup>a</sup> $\pm$ 0.24 | 20.43 <sup>d</sup> $\pm$ 10.30 | 4.00 <sup>b</sup> $\pm$ 0.47 | 1.44 <sup>c</sup> $\pm$ 0.47 | 90.87 <sup>a</sup> $\pm$ 9.25  | 97.97 <sup>a</sup> $\pm$ 9.96  |
| FB-Coll-0038         | 4.63 <sup>a</sup> $\pm$ 1.24 | 1.87 <sup>a</sup> $\pm$ 0.96 | 35.49 <sup>c</sup> $\pm$ 11.45 | 2.74 <sup>c</sup> $\pm$ 0.16 | 2.25 <sup>a</sup> $\pm$ 0.49 | 68.76 <sup>c</sup> $\pm$ 7.98  | 71.74 <sup>c</sup> $\pm$ 10.63 |
| Local check          | 1.82 <sup>c</sup> $\pm$ 0.01 | 1.83 <sup>a</sup> $\pm$ 0.78 | 20.76 <sup>d</sup> $\pm$ 4.65  | 2.00 <sup>e</sup> $\pm$ 0.89 | 2.30 <sup>a</sup> $\pm$ 0.48 | 47.36 <sup>d</sup> $\pm$ 6.89  | 56.48 <sup>c</sup> $\pm$ 5.68  |
| Average              | 3.08 <sup>b</sup> $\pm$ 1.62 | 1.30 <sup>a</sup> $\pm$ 0.68 | 30.94 <sup>c</sup> $\pm$ 19.98 | 2.96 <sup>c</sup> $\pm$ 1.17 | 2.09 <sup>b</sup> $\pm$ 0.49 | 71.37 <sup>c</sup> $\pm$ 11.30 | 77.67 <sup>b</sup> $\pm$ 10.24 |

Means within a column followed by the same letter are not significantly different at 5% probability level, SD = standard deviation, FB = Faba Bean

Table 8. Average yield parameters of faba bean accessions at different experimental sites during 2019 cropping season.

| Faba bean accessions | Angolelana Tera district     |                               |                               | Sululta district              |                              |                                | Midakegn district              |
|----------------------|------------------------------|-------------------------------|-------------------------------|-------------------------------|------------------------------|--------------------------------|--------------------------------|
|                      | Pod/plant                    | Seeds/pod                     | 100 seeds weight (g)          | Pod/plant                     | Seeds/pod                    | 100 seeds weight (g)           | 100 seeds weight (g)           |
|                      | Mean $\pm$ SD                | Mean $\pm$ SD                 | Mean $\pm$ SD                 | Mean $\pm$ SD                 | Mean $\pm$ SD                | Mean $\pm$ SD                  | Mean $\pm$ SD                  |
| FB-Obse              | 0.18 <sup>c</sup> $\pm$ 0.01 | 0.72 <sup>c</sup> $\pm$ 0.02  | 28.38 <sup>b</sup> $\pm$ 0.04 | 7.85 <sup>b</sup> $\pm$ 0.23  | 3.28 <sup>a</sup> $\pm$ 0.25 | 78.69 <sup>a</sup> $\pm$ 2.36  | 73.01 <sup>a</sup> $\pm$ 5.63  |
| FB-28770             | 0.05 <sup>c</sup> $\pm$ 0.02 | 7.30 <sup>a</sup> $\pm$ 0.01  | 9.56 <sup>c</sup> $\pm$ 0.01  | 13.40 <sup>a</sup> $\pm$ 1.23 | 2.71 <sup>a</sup> $\pm$ 0.36 | 25.79 <sup>c</sup> $\pm$ 2.45  | 17.94 <sup>d</sup> $\pm$ 2.35  |
| FB-28107             | 0.41 <sup>c</sup> $\pm$ 0.03 | 2.72 <sup>b</sup> $\pm$ 0.01  | 17.90 <sup>b</sup> $\pm$ 0.04 | 13.39 <sup>a</sup> $\pm$ 1.56 | 2.67 <sup>a</sup> $\pm$ 0.12 | 58.12 <sup>b</sup> $\pm$ 3.21  | 33.51 <sup>c</sup> $\pm$ 2.54  |
| FB-Hachalu           | 0.18 <sup>c</sup> $\pm$ 0.01 | 2.20 <sup>b</sup> $\pm$ 0.03  | 43.40 <sup>a</sup> $\pm$ 0.23 | 5.68 <sup>b</sup> $\pm$ 0.58  | 1.93 <sup>a</sup> $\pm$ 0.47 | 71.39 <sup>a</sup> $\pm$ 1.65  | 75.01 <sup>a</sup> $\pm$ 5.68  |
| FB-26875             | 4.09 <sup>a</sup> $\pm$ 0.02 | 11.02 <sup>a</sup> $\pm$ 0.02 | 9.32 <sup>c</sup> $\pm$ 0.15  | 4.50 <sup>b</sup> $\pm$ 0.24  | 3.11 <sup>a</sup> $\pm$ 0.65 | 21.07 <sup>c</sup> $\pm$ 2.34  | 27.27 <sup>c</sup> $\pm$ 2.45  |
| FB-26873             | 0.54 <sup>c</sup> $\pm$ 0.03 | 0.20 <sup>c</sup> $\pm$ 0.04  | 5.83 <sup>c</sup> $\pm$ 0.12  | 10.84 <sup>a</sup> $\pm$ 1.24 | 2.47 <sup>a</sup> $\pm$ 0.47 | 62.03 <sup>b</sup> $\pm$ 4.35  | 46.20 <sup>b</sup> $\pm$ 3.65  |
| FB-26883             | 0.41 <sup>c</sup> $\pm$ 0.04 | 0.48 <sup>c</sup> $\pm$ 0.01  | 16.71 <sup>b</sup> $\pm$ 1.30 | 4.90 <sup>b</sup> $\pm$ 0.45  | 3.12 <sup>a</sup> $\pm$ 0.36 | 23.53 <sup>c</sup> $\pm$ 2.35  | 49.20 <sup>b</sup> $\pm$ 2.45  |
| FB-Numan             | 5.31 <sup>a</sup> $\pm$ 0.01 | 10.4 <sup>a</sup> $\pm$ 0.06  | 41.01 <sup>a</sup> $\pm$ 2.16 | 14.18 <sup>a</sup> $\pm$ 2.69 | 1.39 <sup>a</sup> $\pm$ 0.21 | 60.34 <sup>b</sup> $\pm$ 2.64  | 66.13 <sup>a</sup> $\pm$ 3.65  |
| FB-26867             | 0.32 <sup>c</sup> $\pm$ 0.04 | 0.32 <sup>c</sup> $\pm$ 0.01  | 4.50 <sup>d</sup> $\pm$ 0.69  | 10.57 <sup>a</sup> $\pm$ 2.35 | 1.92 <sup>a</sup> $\pm$ 0.84 | 50.33 <sup>b</sup> $\pm$ 2.65  | 54.27 <sup>b</sup> $\pm$ 4.56  |
| FB-Tumsa             | 1.64 <sup>b</sup> $\pm$ 0.06 | 2.51 <sup>b</sup> $\pm$ 0.04  | 55.34 <sup>a</sup> $\pm$ 2.63 | 6.56 <sup>b</sup> $\pm$ 1.54  | 1.79 <sup>a</sup> $\pm$ 0.65 | 31.08 <sup>c</sup> $\pm$ 2.35  | 70.71 <sup>a</sup> $\pm$ 4.65  |
| FB-26885             | 4.95 <sup>a</sup> $\pm$ 0.04 | 9.13 <sup>a</sup> $\pm$ 0.08  | 37.60 <sup>a</sup> $\pm$ 1.56 | 8.32 <sup>b</sup> $\pm$ 0.85  | 2.40 <sup>a</sup> $\pm$ 0.14 | 32.32 <sup>c</sup> $\pm$ 2.45  | 6.88 <sup>d</sup> $\pm$ 1.25   |
| FB-25280             | 0.00 <sup>d</sup> $\pm$ 0.00 | 0.00 <sup>c</sup> $\pm$ 0.00  | 0.00 <sup>d</sup> $\pm$ 0.00  | 11.53 <sup>a</sup> $\pm$ 1.45 | 2.01 <sup>a</sup> $\pm$ 0.90 | 23.65 <sup>c</sup> $\pm$ 2.34  | 23.12 <sup>c</sup> $\pm$ 2.35  |
| FB-26872             | 1.09 <sup>b</sup> $\pm$ 0.03 | 2.49 <sup>b</sup> $\pm$ 0.05  | 7.97 <sup>c</sup> $\pm$ 1.25  | 6.88 <sup>b</sup> $\pm$ 0.24  | 2.48 <sup>a</sup> $\pm$ 0.21 | 63.71 <sup>b</sup> $\pm$ 2.68  | 47.01 <sup>b</sup> $\pm$ 5.68  |
| FB- Coll-0038        | 3.45 <sup>a</sup> $\pm$ 0.01 | 3.91 <sup>b</sup> $\pm$ 0.03  | 13.85 <sup>c</sup> $\pm$ 2.03 | 9.92 <sup>b</sup> $\pm$ 0.69  | 3.09 <sup>a</sup> $\pm$ 0.47 | 34.16 <sup>c</sup> $\pm$ 4.68  | 46.26 <sup>b</sup> $\pm$ 6.35  |
| Local check          | 0.60 <sup>c</sup> $\pm$ 0.03 | 2.33 <sup>b</sup> $\pm$ 0.01  | 7.82 <sup>c</sup> $\pm$ 1.06  | 10.36 <sup>a</sup> $\pm$ 2.56 | 1.98 <sup>a</sup> $\pm$ 0.25 | 48.08 <sup>b</sup> $\pm$ 4.52  | 39.52 <sup>b</sup> $\pm$ 4.56  |
| Average              | 1.55 <sup>b</sup> $\pm$ 1.09 | 3.72 <sup>b</sup> $\pm$ 0.83  | 19.95 <sup>b</sup> $\pm$ 6.96 | 9.26 <sup>b</sup> $\pm$ 3.16  | 2.42 <sup>a</sup> $\pm$ 0.57 | 45.62 <sup>b</sup> $\pm$ 13.60 | 45.07 <sup>b</sup> $\pm$ 10.78 |

Means within a column followed by the same letter are not significantly different at 5 % probability level, SD = standard deviation, FB = Faba Bean

### **Analysis of yield parameters during 2018–2019 cropping seasons**

The interaction of faba bean accessions with gall disease significantly ( $p < 0.05$ ) affected yield parameters over districts. The number of pods/plants was higher at Sululta (6.11 pods/plant) than Angolelana Tera (2.18 pods/plant) district. However, there was no significant difference ( $p > 0.05$ ) among the districts on the number of seeds/pods in a combined analysis of accessions (Table 9). Lower (5.73 g) 100 seeds weight was recorded at Angolelana Tera than other districts (Table 9).

Maximum pods per plant were recorded on FB-26875, and FB-Numan accessions at Angolelana Tera, and Sululta districts, respectively. There was no significant ( $p > 0.05$ ) difference recorded among evaluated accessions in many seeds per pod at Sululta district, while FB-Coll-0038 (2.67 seeds/pod), FB-Obse (2.64 seeds/pod), and FB-26867 (2.46 seeds/pod) ranked first to third for the number of seeds per pod. There was a significant difference ( $p < 0.05$ ) among accessions in 100 seeds weight at both districts. The maximum 100 seeds weight was recorded on FB-Obse (78.69 g) at Sululta followed by FB-Hachalu (75 g), and FB-Numan (12 g) at Midakegn and Angolelana Tera districts, respectively (Table 9).

In combined mean analysis over the cropping seasons (2018–2019), FB-Obse, FB-26872, FB-26873, and FB-Hachalu produced significantly ( $p < 0.05$ ) better 100 seeds weight than others with the value 82.50 g, 77.29 g, 71.08 g, and 71.32 g at Sululta, respectively. At Midakegn, FB-Obse (81.91 g), FB-Tumsa (79.78 g), FB-Hachalu (79.66 g), and FB-26872 (78.49 g) exhibited the highest 100 seeds weight, whereas FB-Numan (56.77 g), FB-26885 (51.25 g), FB-Tumsa (45.76 g), and FB-Obse (41.26 g) resulted in the highest 100 seeds weight among others at Angolelana Tera district. In general, an average maximum 100 seeds weight was harvested from Midakegn (61.41 g) followed by Sululta (58.50 g) and Angolelana Tera (25.44 g) districts (Table 9).

Table 9. Average yield parameters of faba bean accessions during 2018–2019 cropping seasons.

| Faba bean accessions | Angolelana Tera district |               |                      | Sululta district |                |                      | Midakegn district    |
|----------------------|--------------------------|---------------|----------------------|------------------|----------------|----------------------|----------------------|
|                      | Pod/plant                | Seeds/pod     | 100 seeds weight (g) | Pod/plant        | Seeds/pod      | 100 seeds weight (g) | 100 seeds weight (g) |
|                      | Mean ± SD                | Mean ± SD     | Mean ± SD            | Mean ± SD        | Mean ± SD      | Mean ± SD            | Mean ± SD            |
| FB-Obse              | 1.15c* ± 4.08            | 0.51b* ± 1.00 | 41.26a* ± 0.06       | 4.85a** ± 3.06   | 2.64a** ± 0.00 | 82.50d** ± 11.25     | 81.91d** ± 2.01      |
| FB-28770             | 2.89b* ± 1.05            | 4.51a* ± 1.13 | 15.00a* ± 0.07       | 8.20b** ± 1.03   | 2.17a** ± 0.04 | 45.56b** ± 4.07      | 44.00b** ± 3.00      |
| FB-28107             | 1.15c* ± 1.10            | 1.73c* ± 0.50 | 26.17a* ± 0.70       | 8.01b** ± 1.20   | 2.39a* ± 0.10  | 63.67a** ± 6.60      | 58.05a** ± 3.01      |
| FB-Hachalu           | 2.00d* ± 0.01            | 2.00d* ± 0.10 | 40.00c* ± 0.40       | 3.74a** ± 2.19   | 2.33a** ± 0.80 | 71.32d** ± 6.02      | 79.66d** ± 3.00      |
| FB-26875             | 4.08a* ± 1.33            | 6.50a* ± 2.29 | 15.00a* ± 0.67       | 4.35a* ± 0.11    | 2.26a** ± 0.26 | 51.90a** ± 19.07     | 54.00a** ± 3.07      |
| FB-26873             | 1.15c* ± 3.01            | 0.77b* ± 1.12 | 10.11b* ± 0.14       | 6.93a** ± 2.12   | 2.12a** ± 0.54 | 71.08a** ± 7.01      | 62.62c** ± 1.00      |
| FB-26883             | 3.00b* ± 3.01            | 0.58b* ± 0.06 | 17.19a* ± 0.70       | 4.27a** ± 2.09   | 2.39a** ± 0.05 | 45.24b** ± 3.03      | 61.00c** ± 1.00      |
| FB-Numan             | 4.00a* ± 2.80            | 6.00a* ± 0.15 | 56.77b* ± 0.18       | 8.28b** ± 0.71   | 1.96a** ± 0.12 | 62.99c** ± 14.15     | 69.00d** ± 2.00      |
| FB-26867             | 1.00c* ± 3.07            | 0.29b* ± 1.06 | 10.00b* ± 1.09       | 6.35a** ± 3.15   | 2.46a** ± 0.19 | 64.97c** ± 10.25     | 63.00c** ± 1.00      |
| FB-Tumsa             | 2.89b* ± 4.14            | 2.02c* ± 0.07 | 45.76a* ± 0.15       | 4.13a** ± 2.10   | 2.28a* ± 0.26  | 50.54a** ± 2.15      | 79.78d*** ± 3.00     |
| FB-26885             | 3.61b* ± 1.03            | 5.51a* ± 2.22 | 51.25a* ± 1.12       | 5.80a** ± 3.34   | 2.15a** ± 0.10 | 44.24b** ± 1.40      | 40.00b** ± 1.00      |
| FB-25280             | 0.00d* ± 0.00            | 0.00d* ± 0.00 | 0.00c* ± 0.00        | 8.80b** ± 3.06   | 1.96a** ± 0.08 | 46.96b** ± 0.07      | 48.00a** ± 2.00      |
| FB-26872             | 2.52b* ± 1.01            | 2.08c* ± 0.50 | 14.20a* ± 0.20       | 5.44a** ± 3.50   | 1.96a* ± 0.29  | 77.29c** ± 24.20     | 72.49c** ± 1.02      |
| FB- Coll-0038        | 4.04a* ± 2.50            | 2.89c* ± 1.06 | 24.67a* ± 0.00       | 6.33a* ± 2.14    | 2.67a* ± 0.29  | 51.46a** ± 6.15      | 59.00c** ± 4.00      |
| Local check          | 1.21c* ± 1.12            | 2.08c* ± 0.11 | 14.29a* ± 0.19       | 6.18a** ± 4.12   | 2.14a* ± 0.27  | 47.72b** ± 9.32      | 48.00a** ± 2.00      |
| Average              | 2.18b* ± 1.30            | 2.36c* ± 0.70 | 25.44a* ± 2.27       | 6.11a** ± 1.27   | 2.26a* ± 0.13  | 58.50a** ± 10.27     | 61.41c** ± 2.08      |

Means within a column followed by the same letter and means across the district within a row followed by the same number of \* are not significantly different at 5%, SD = standard deviation, FB = Faba Bean

## P-values of contrast analysis between different traits

The combined mean analysis over cropping seasons, and districts showed that there was significant difference ( $p < 0.05$ ) in disease severity and 100 seeds weight between local check and various faba bean accessions. Gall disease severity and faba bean yield parameters in FB-Obse accession showed no significant difference ( $p > 0.05$ ) from FB-26872 accession. Significant ( $p = 0.003$ ) higher disease severity (DSI = 26.96%) was recorded during 2019 than others cropping seasons (Table 10). On the other hand, significant ( $p < 0.05$ ) higher seeds/pod (2.8 seeds) and 100 seeds weight (55.8 g) yield parameters were recorded in 2018 than 2019 cropping seasons over the three districts. Also, significant variations were recorded in gall disease severity and yield parameters among districts. The highest disease severity was recorded at Angolelana Tera (DSI = 29.26%) among Sululta and Midakegn districts. Maximum 100 seeds weight was recorded at Midakegn district (61.43 g) although it was not significantly different ( $p = 0.445$ ) from Sululta district (Table 10).

Table 10. P-values of contrast analysis between different traits.

| Contrast                                 | DF  | Severity | Pod/Plant | Seed/Pod | 100 seeds weight |
|------------------------------------------|-----|----------|-----------|----------|------------------|
| <b>Between some faba bean accessions</b> |     |          |           |          |                  |
| Local check vs FB-Obse                   | 134 | <0.001*  | 0.778     | 0.962    | 0.004*           |
| FB-Obse vs FB-26872                      | 134 | >0.05    | >0.05     | >0.05    | >0.05            |
| FB-Obse vs FB-26885                      | 134 | 0.007*   | 0.360     | 0.139    | 0.035*           |
| FB-Obse vs FB-26867                      | 134 | 0.164    | 0.820     | 1.000    | 0.041*           |
| <b>Between cropping seasons</b>          |     |          |           |          |                  |
| 2017 vs 2018                             | 134 | 0.437    | NA        | NA       | NA               |
| 2017 vs 2019                             | 134 | 0.023*   | NA        | NA       | NA               |
| 2018 vs 2019                             | 134 | 0.003*   | 0.044     | 0.397    | 0.621            |
| <b>Between districts</b>                 |     |          |           |          |                  |
| Angolelana Tera vs Sululta               | 134 | <0.001*  | <0.001*   | 0.530    | <0.001*          |
| Angolelana Tera vs Midakegn              | 134 | <0.001*  | 0.002*    | 0.185    | <0.001*          |
| Sululta vs Midakegn                      | 134 | <0.001*  | <0.001*   | 0.483    | 0.445            |

DF = degrees of freedom; FB = faba bean; “\*” = significant at 5%, NA = not analyzed

## Correlation analysis between different traits

The Pearson's correlation coefficient showed that gall disease incidence has weak, positive and high significant ( $p < 0.05$ ) correlation with disease severity index ( $r = 0.49$ ) (Table 11). Likewise, a weak, positive and significant correlation of seeds per pod with 100 seeds weight ( $r = 0.06$ ) was observed. Similarly, positive and significant correlation of altitude ( $r = 0.27$ ) and precipitation ( $r = 0.37$ ) with gall disease severity was also observed. In contrary, pods per plant ( $r = -0.05$ ), seeds per pod ( $r = -0.03$ ), and 100 seeds weight ( $r = -0.14$ ) was negatively correlated with gall disease

incidence. Also, pods per plant ( $r = -0.07$ ), 100 seeds weight ( $r = -0.53$ ), and temperature ( $r = -0.23$ ) have significant negative correlation with gall disease severity index (Table 11).

Table 11. Correlation coefficients values of the pairs of different traits.

| Traits code | Traits           | T1      | T2      | T3      | T4      | T5     | T6     | T7      | T8      | T9 |
|-------------|------------------|---------|---------|---------|---------|--------|--------|---------|---------|----|
| T1          | FB-accession     | 1       |         |         |         |        |        |         |         |    |
| T2          | Incidence        | 0.06*   | 1       |         |         |        |        |         |         |    |
| T3          | Disease severity | 0.12**  | 0.49**  | 1       |         |        |        |         |         |    |
| T4          | Pods per plant   | 0.08**  | -0.05ns | -0.07** | 1       |        |        |         |         |    |
| T5          | Seeds per pod    | 0.05*   | -0.03ns | -0.02ns | 0.08**  | 1      |        |         |         |    |
| T6          | seeds weight     | 0.18**  | -0.14** | -0.53** | 0.04ns  | 0.06*  | 1      |         |         |    |
| T7          | Altitude         | 0.01ns  | 0.49**  | 0.27**  | 0.31**  | 0.05ns | 0.10** | 1       |         |    |
| T8          | Precipitation    | 0.01ns  | 0.64**  | 0.37**  | 0.28**  | 0.04ns | 0.01** | 0.94**  | 1       |    |
| T9          | Temperature      | -0.01ns | -0.43** | -0.23** | -0.31** | 0.08** | 0.10** | -0.99** | -0.89** | 1  |

\*\*\*: Correlation is significant at the 0.01 level (2-tailed)

\*: Correlation is significant at the 0.05 level (2-tailed), ns: not significant, FB: Faba bean

## Regression analysis

The simple linear regression analysis revealed that for each unit increase of an altitude resulted in a raise of gall disease incidence by 0.16% (Fig. 3A) and disease severity index by 0.09% (Fig. 3B). Based on the coefficient of determination value ( $R^2$ ), the equation explained about 92% of the variation in disease incidence and 98% of the variation in gall disease severity index were due to altitude. Similarly, the estimated slope of the regression line obtained for gall disease incidence was 0.43. The estimate showed that for each unit increase in gall disease incidence, there was a raise in gall disease severity index by 0.43%. The equation explained about 88% of the variation in severity index due to disease incidence (Fig. 3C). Also, a unit increase of pods per plant resulted in an increase of seeds per pod by 0.47 seeds with 96% variations (Fig. 3E). On the contrary, for each unit increase in percent severity index of gall disease, there was a 100 seeds weight loss by 0.19 g. The regression equation explained about 99% variation in seed weight loss by disease severity (Fig. 3D). For each unit increase of percent severity index of gall disease there was pods loss by 0.045 pods (Fig. 3F).




Fig. 3. Regression analysis: A. Altitude and percent disease incidence, B. Altitude and Percent severity index, C. Percent disease incidence and severity index, D. Percent severity index and 100 seeds weight, E. Pods/plant and seeds/pod, F. Percent severity index and pods/plant.



## DISCUSSION

Identification of faba bean accessions resistant to gall disease with a high yield potential could guarantee the use of eco-friendly disease management options and cultivation productivity for faba bean farmers. Resistant accessions reside in high value among disease controlling methods, because it is certainly applied, requires few inputs and is economically profitable (Behailu Mulugeta *et al.*, 2018). The gall disease was characterized by the formation of chlorotic galls which enlarges to light brown, circular, or elliptical rough spots on both sides of the leaves and stems (Teklay Abebe *et al.*, 2014). The disease was observed soon after crop emergence on infested soil and faba bean residue plots. It is more destructive at the early growth stages of the plant, is spread by wind, rain, and weeding practices, and causes a high reduction of crop yield (Teklay Abebe *et al.*, 2018).

Interestingly, we observed similar disease symptoms with the previous reports in our study (Teklay Abebe *et al.*, 2014). However, the date of occurrence of gall disease after the crop emergence varied at different experimental sites and cropping seasons. The progress of the disease incidence gradually became severe at high altitude and average low temperature as the rain increases and reaches its maximum in the mid rainy season. These could be correlated with the climate of the experimental locations and cropping seasons. Pszczółkowska *et al.* (2020) reported that disease incidence and spread of pathogenic fungi were related to weather conditions and determined by the distribution pattern of precipitation and average temperature during the growing season. Besides, the distribution of gall disease has a linear relationship with the rise of altitude (Teklay Abebe *et al.*, 2014).

In the present study, the absence of innate resistance was observed as no faba bean variety was found resistant to gall disease. However, FB-26872, FB-Tumsa, FB-26873, FB-28107, FB-Obse, FB-Hachalu, FB-26883, FB-Numan, FB-28770, FB-26885, FB-COLL-0038, FB-26875, FB-26867, and FB-25280 accessions had shown better reactions against gall disease. The reaction levels varied at different experimental locations and during main cropping seasons. They demonstrated better seed field germination percentage, lower disease severity, AUDPC, and apparent infection rate against gall disease causal agent *Olpidium viciae* compared with the rest of the accessions.

These results agree with Teklay Abebe *et al.* (2018), who reported that improved and local faba bean varieties were affected by faba bean gall

disease without any tolerance. According to Bogale Nigir *et al.* (2017), gall disease infected both local cultivars and improved varieties in Ethiopia; nevertheless, the intensities were higher on the local cultivars. This suggests that the difference in reaction levels and disease occurrence rate among varieties could be associated with genotypic variability in the inheritance of gall disease resistant traits of the crops (Sheikh *et al.*, 2015; Méndez-Natera *et al.*, 2016; Behailu Mulugeta *et al.*, 2018), a variation of the previous inoculum level of the disease pathogen (Teklay Abebe *et al.*, 2014), soil types (Endalkachew Fekadu *et al.*, 2018), and climate of the study area (Pszczółkowska *et al.*, 2020). The variation of the climate could significantly influence the infection rate and spread of conidia of the test pathogen (Teklay Abebe *et al.*, 2014). Significant variability among varieties allows us to select the faba bean accessions with better tolerance against gall disease.

In the present study, fourteen faba bean accessions revealed high yield performance among the tested accessions and local check at each experimental site and cropping seasons. A maximum of 100 seeds weight was harvested in the 2018 cropping season. Likewise, a maximum of 100 seeds weight was harvested from Midakegn, followed by Sululta, and Angolelana Tera districts. These findings were in line with Endalkachew Fekadu *et al.* (2018) who reported on the yield performance of faba bean varieties in Ethiopia in the case of pods per plant, and 100 seed weight, while slightly contradicted in the case of the number of seeds per pod. In agreement with the current finding, Bogale Nigir *et al.* (2017) stated that the number of pods per plant was significantly affected by faba bean varieties. Basha Kebede and Dembi Korji (2017) also reported a significant difference among faba bean varieties on yield parameters.

The observed difference of yield parameters could be correlated with the climate of the cropping seasons, inherent genetic differences of faba bean varieties, variation in the adaptive ability of the crops to the study area, the severity level of gall disease, and the occurrence of frost (Basha Kebede and Dembi Korji, 2017). The disease resistant variety can boost crop yield (Méndez-Natera *et al.*, 2016). The climate of the study location influenced the growth and yielding of faba beans (Amalfitano *et al.*, 2018; Pszczółkowska *et al.*, 2020).

This study confirmed by Pearson correlation coefficient and regression analysis result the impact of climate of the study sites on faba bean gall disease pressure and yield parameters. The Pearson correlation coefficient

implied that altitude, precipitation, and average low temperature positively affected gall disease severity, while gall disease severity negatively influenced yield parameters. The linear regression analysis showed that for each unit increase in altitude of the study sites, there was an increment of percent disease severity index by 0.09%. This accounts for 98% variation in the faba bean gall disease percent severity index. On the other hand, for each unit increase in the percent severity index, 100 seeds weight of faba bean accessions decreased by 0.19% and accounted for 99% variations in 100 seeds weight of the crops.

### CONCLUSIONS AND RECOMMENDATIONS

Faba bean accessions responded at various levels against gall disease at different study areas and cropping years. Gall disease negatively influenced faba bean yield parameters. No faba bean accessions was found resistant to gall disease. FB-Obse, FB-Hachalu, FB-26873, FB-Numan, and FB-26872 accessions revealed low DSI and high crop yield. However, they exhibited various level response in different study areas. Thus, a single faba bean variety cannot be considered best fit for all study sites. The following accessions; FB-Obse, FB-26872, FB-26873, and FB-Hachalu at Sululta; FB-Obse, FB-Tumsa, FB-Hachalu, and FB-26872 at Midakegn; whereas FB-Numan, FB-26885, FB-Tumsa, and FB-Obse at Angolelana Tera district are promising to the corresponding localities to be used for the management of faba bean gall disease. Efforts have to be made by researchers to develop faba bean varieties with gall disease resistant lines at multiple faba bean production areas.

### ACKNOWLEDGEMENTS

We would like to express our sincere gratitude to Addis Ababa University Office for Research, Thematic Research Directorate for funding this research. We are profoundly grateful to the Ethiopian Biodiversity Institute (EBI) and Holeta Agricultural Research Centre (HARC) for the provision of various faba bean accessions.

### REFERENCES

- Aghajani, M.A. (2009). Stemphylium leaf blight of broad bean in Iran. *J. Plant Pathol.* **91**: 97–112.
- Alireza, N., Afshin, S., Ali, R., Amir, D. and Faranak, N. (2021). Determination of soybean yield gap and potential production in Iran using modeling approach and GIS. *J. Integr. Agric.* **20**: 395–407.
- Amalfitano, C., Agrelli, D., Borrelli, C., Cuciniello, A., Morano, G. and Caruso, G. (2018). Production system effects on growth, pod yield and seed quality of organic faba bean in southern Italy. *Folia Hortic.* **30**: 375–385.

- Basha Kebede and Dembi Korji (2017). Pre-scaling up of improved faba bean technologies in the highland districts of Guji Zone, Oromia regional state, Ethiopia. *Asian J. Agric. Rural Dev.* **7**: 115–119.
- Behailu Mulugeta, Kifle Dagne, Gemechu Keneni, Seid Ahmed and Kassahun Tesfaye (2018). Genetic diversity study of Ethiopian faba bean (*Vicia faba* L.) varieties based on phenotypic traits and inter simple sequence repeat (ISSR) markers. *Afr. J. Biotechnol.* **17**: 433–446.
- Bhagat, S., Bambawale, O., Tripathi, A., Ahmad, I. and Srivastava, R. (2013). Biological management of fusarial wilt of tomato by *Trichoderma* spp. in Andamans. *Indian J. Hortic.* **70**: 397–403.
- Bitew Beyene and Tigabie Abiro (2016). Management of faba bean gall disease (Kormid) in North Shewa highlands, Ethiopia. *Adv. Crop Sci. Technol.* **4**(4): 225.
- Bogale Nigir, Mulugeta Tilahun, Setu Bazie, Tesifaye Desale and Eyeberu Abere (2017). Management of faba bean gall in faba bean producing area of Eastern Amhara, Ethiopia. *J. Agric. Environ. Int. Dev.* **111**: 343–350.
- EMA (Ethiopian Meteorology Agency) (2019). Monthly climate data of the year 2017–2019 at Angolelana Tera, Sululta, and Midakegn districts. Addis Ababa, Ethiopia.
- Endalkachew Fekadu, Kibebew Kibret, Asmare Melese and Bobe Bedadi (2018). Yield of faba bean (*Vicia faba* L.) as affected by lime, mineral P, farmyard manure, compost and rhizobium in acid soil of Lay Gayint District, northwestern highlands of Ethiopia. *Agric. Food Secur.* **7**: 16.
- Etemadi, F., Hashemi, M., Zandvakili, O. and Liu, X. (2019). Agronomy, nutritional value, and medicinal application of faba bean (*Vicia faba* L.). *Hortic. Plant J.* **5**: 170–182.
- Méndez-Natera, J.R., Luna-Tineo, J.A., Barrios-Azócar, L.A. and Cedeño, J.R. (2016). Screening of Indian peanut genotypes for resistance to *Cercospora* leaf-spot under savanna conditions. *Emir. J. Food Agric.* **28**: 833–841.
- Pornthip, R. and Prakrit, S. (2021). Identification and resistant characterization of legume sources against *Meloidogyne incognita*. *J. Integr. Agric.* **20**: 168–177.
- Pszczółkowska, A., Okorski, A., Fordoński, G., Kozak, M. and Dzienis, G. (2020). Effect of weather conditions on yield and health status of faba bean seeds in Poland. *Agron.* **10**: 48.
- Sheikh, F., Dehghani, H. and Aghajani, A. (2015). Screening *Vicia faba* L. genotypes for resistance to *Stemphylium* blight in Iran. *Eur. J. Plant Pathol.* **143**: 677–689.
- Singh, A., Bharati, R., Naresh, M. and Anitha, P. (2013). An assessment of faba bean (*Vicia faba* L.) current status and future prospect. *Afr. J. Agric. Res.* **8**: 6634–6641.
- Soler, A., Marie-Alphonsine, P., Qu'en'éherv'e, P., Prin, Y., Sanguin, H., et al. (2021). Field management of *Rotylenchulus reniformis* on pineapple combining crop rotation, chemical mediated induced resistance and endophytic bacterial inoculation. *Crop Prot.* **141**: 105–446.
- Su, Z., Liu, Z., Bai, F., Zhang, Z., Sun, S., Huang, Q., Liu, T., Liu, X. and Yang, X. (2021). Cultivar selection can increase yield potential and resource use efficiency of spring maize to adapt to climate change in Northeast China. *J. Integr. Agric.* **20**: 371–382.
- Teklay Abebe, Gurja Belay, Gemechu Keneni, Taye Tadesse (2018). Fungicidal management of the newly emerging faba bean disease “gall” (*Olpidium viciae* Kusano) in Tigray, Ethiopia. *Crop Prot.* **107**: 19–25.
- Teklay Abebe, Tsehay Birhane, Yemane Nega and Assefa Workineh (2014). Study on

occurrence and importance of faba bean disease with special consideration to the newly emerging “faba bean gall” in Tigray, Ethiopia. *Afr. J. Agric. Res.* **9**: 3627–3631.

Villegas-Fernandez, M., Sillero, C. and Rubiales, D. (2011). Screening faba bean for chocolate spot resistance: evaluation methods and effects of age of host tissue and temperature. *Eur. J. Plant Pathol.* **132**: 443–453.

Wondafrash Mulugeta, Kindie Tesfaye, Mezegebu Getnet, Seid Ahmed, Amsalu Nebiyu, and Fasil Mekuanint (2019). Quantifying yield potential and yield gaps of faba bean in Ethiopia. *Ethiop. J. Agric. Sci.* **29**(3): 105–120.